

UTILITY PATENT APPLICATION TRANSMITTAL

DUPLICATE

Address to: Commissioner for Patents Box PATENT APPLICATION Washington, DC 20231	Attorney Docket No. REF/FOSSHEIM/100
First Named Inventor (or Identifier) FOSSHEIM et al.	Total Pages 112

Transmitted herewith is a patent application under 37 CFR 1.53(b).

Entitled: **USE OF PARTICULATE CONTRAST AGENTS IN DIAGNOSTIC IMAGING FOR STUDYING PHYSIOLOGICAL PARAMETERS**

- ☒ 1. Submitted herewith are the following:
- 79 pages of specification.
 - ☒ Abstract.
 - 13 sheet(s) of drawings.
 - 22 claim(s).
 - ☒ Oath/Declaration signed by each inventor.
 - ☐ signed Inventor Small Entity Statement(s).
 - ☐ signed non-Inventor Small Entity Statement(s).
 - ☐ signed Small Business Small Entity Statement(s).
 - ☐ signed Non-Profit Small Entity Statement(s).
 - ☒ Preliminary Amendment.
 - ☒ Information Disclosure Statement(s).
 - 1 page of Form PTO-1449.
 - ☒ Assignment of the invention, Cover Sheet, and payment of the \$40.00 recordal fee.
 - ☐ certified copy of application no. _____ filed in _____. Priority is claimed.
 - ☒ check in the amount of \$ 786.00 including any assignment recordal fee.
- ☒ 2. The Commissioner is authorized to credit any overpayment and charge any deficiency in any fees required under 37 CFR 1.16 and/or 1.17, to Deposit Account No. 02-0200.
- ☐ 3. Insert before the first sentence of the specification: - - This application claims the benefit of provisional application number _____ filed _____ - -
- ☐ 4. Insert before the first sentence of the specification: - - This application is a Continuation-in-part of nonprovisional application number _____ filed _____ - -
- ☐ 5. Other: _____

THE FILING FEE IS CALCULATED AS FOLLOWS:

THE FILING FEE IS CALCULATED AS FOLLOWS:				Basic Fee*	\$710 00
Total Claims:	22	- 20 =	2	X \$18 =	36.00
Independent Claims:	3	- 3 =	0	X \$80 =	0 00
Correspondence Address BACON & THOMAS, PLLC 625 Slaters Lane, 4 th Floor Alexandria, VA 22314-1176				Multiple Dependent Claim (add \$270.00):	
				Subtotal:	746 00
				50% Reduction if Small Entity Status:	
Phone: 703-683-0500 Fax: 703-683-1080				Total:	746.00

Date: October 6, 2000	Name: Richard E. Fichter	Signature: <i>Richard E. Fichter</i>	Reg. No. 26,382
--------------------------	-----------------------------	---	--------------------

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re continuation application of: :
FOSSHEIM et al. : Office of Initial Patent Examination
Continuation of PCT/GB99/01100 :
This application filed: October 6, 2000 :
For: Use of Particulate Contrast Agents in Diagnostic Imaging for Studying :
Physiological Parameters :

**PRELIMINARY AMENDMENT
AND INFORMATION DISCLOSURE STATEMENT**

Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

Please amend the above-identified application as follows:

IN THE SPECIFICATION:

Please insert as the first sentence - -This application is a continuation of PCT/GB99/01100, filed April 9, 1999 (of which the entire disclosure of the pending, prior application is hereby incorporated by reference), which itself is a continuation-in-part of application number 60/119,808, filed February 12, 1999.- -

Please add the attached Abstract of the Disclosure to the application.

IN THE CLAIMS:

Claim 4, line 1, please cancel "any one of claims 1 to 3" and insert - -claim 1- -.

Claim 5, line 1, please cancel "any one of claims 1 to 4" and insert - -claim 1- -.

Claim 12, line 1, please cancel "or 11".

Claim 15, lines 1 and 2, please cancel "any one of the preceding claims" and insert - -claim 1- -.

Claim 16, line 1, please cancel "any one of claims 1 to 14" and insert - -claim 1- -.

Claim 17, line 1, please cancel "any one of claims 1 to 16" and insert - -claim 1- -.

Please cancel claim 22 without prejudice or disclaimer.

REMARKS

Applicants have amended the specification to cross reference the parent applications and to add an Abstract of the Disclosure to the application.


Applicants have amended the claims in order to reduce the initial filing fee by deleting the improper use claim 22 and to delete the multiple dependent claims from the application. Applicants retain the right to reintroduce any subject matter canceled by the present Amendment at any time during the prosecution of this application or any further application claiming benefit of this application.

Applicants are submitting herewith a copy of the Search Report which issued on International Application No. PCT/GB99/01100, of which the present application is a continuation. All of the publications cited in the International Search Report are listed on the attached Form PTO-1449.

In view of the above amendments to the claims, an early action on the application is most respectfully requested.

Respectfully submitted,

BACON & THOMAS, PLLC

By: 
Richard E. Fichter
Registration No. 26,382

625 Slaters Lane, 4th Fl.
Alexandria, Virginia 22314
Phone: (703) 683-0500
Facsimile: (703) 683-1080

REF:kdd
PA01 wpd

October 6, 2000

USE OF PARTICULATE CONTRAST AGENTS IN DIAGNOSTIC IMAGING FOR STUDYING PHYSIOLOGICAL PARAMETERS

This invention relates to the use of particulate contrast agents in diagnostic imaging procedures for studying physiological parameters of the subject under investigation.

In diagnostic imaging procedures, e.g. X-ray, MRI, ultrasound, light imaging and nuclear imaging, it has long been known to use contrast agents to facilitate visualization of particular organs or tissues or to identify diseased or malfunctioning regions, ie. generating morphological images.

The present invention is concerned with the use of parenterally administered particulate contrast agents for the quantitative or qualitative study of physiological parameters within the human or non-animal (e.g. mammalian, avian or reptilian, but preferably mammalian) body.

Such parameters include for example pH, temperature, pressure, oxygen tension, carbon dioxide tension, ion tension/concentration the presence or concentration of other body metabolites or enzymes and cell surface properties, e.g. the presence or absence of various cell surface receptors. Parameters such as these may be indicative of the normal or abnormal functioning of the body as a whole or of a particular localized region, e.g. an organ which may or may not be tumorous, infected or otherwise malfunctioning. Likewise variations in such parameters may occur in response to drugs or other treatments administered to the body, e.g. hyperthermic treatment. As a result, quantitative, semi-quantitative or even qualitative determination of such parameters may be used to assess the need for a particular treatment or to monitor the success of a particular treatment.

pH and temperature are particularly important as

- 2 -

indicators of abnormality or malfunction.

Several in vivo methods, both imaging techniques and non-imaging techniques, can be used to study physiological parameters, e.g. to diagnose disease. Typical non-imaging techniques include simple blood pressure measurements, electrocardiography or electrocephalography for detection of electric currents in the heart muscle and brain, respectively, and other simple tests performed in doctors' offices or hospitals. Today, a variety of imaging techniques are also used. The most frequently used methods include various X-ray based techniques, MRI, ultrasound and diagnostic methods based on radioactive materials (e.g. scintigraphy, PET and SPECT). Other diagnostic imaging methods include light imaging modalities, Overhauser MR (OMRI), oxygen imaging (OXI) which is based on OMRI, magnetic source imaging (MSI), applied potential tomography (APT) and imaging methods based on microwaves.

The images obtained in X-ray techniques reflect the different densities of structures/organs/tissues in the patient's body. Contrast agents are today used to improve the image contrast in soft tissue examinations. Examples of such contrast agents include gas (negative contrast effect relative to tissue); barium sulphate suspensions; and iodinated agents including ionic monomeric agents, non-ionic monomers, ionic dimers and non-ionic dimers. Typical examples of commercial X-ray contrast agents are Omnipaque® and Visipaque®.

MRI is an imaging method generally based on interactions between radiowaves and body tissue water protons in a magnetic field. The contrast parameter or signal intensity is dependent on several factors including proton density, spin lattice (T_1) and spin spin (T_2) relaxation times of water protons. Typical commercial MRI contrast agents include Omniscan®, Magnevist® and ProHance®.

Ultrasound is another valuable modality in

- 3 -

diagnostic imaging as it does not involve the use of ionizing radiation. In ultrasound examinations the patient is generally exposed to sound waves in the frequency of 1-10 MHz. These sound waves (or ultrasound waves) penetrate through or are reflected from the tissue. The transmitted or reflected sound waves are detected by a "microphone" and form the basis for development of a ultrasound image. Ultrasound imaging is a method of choice in pregnancy checks and birth control and diagnosis of cardiovascular and liver diseases.

Although ultrasound contrast agents have been approved, there is as yet no general use of these agents. The main reason for this is the poor efficacy of the "first generation" agents. The ultrasound contrast agents currently under development are based on encapsulated gas because the reflection of sound from the liquid-gas interface is extremely efficient.

Typical ultrasound contrast agents are gas encapsulated in a sugar matrix, in a shell of denaturated albumin/or partly denaturated albumin, in polymers, and in surfactants including phospholipids. A typical ultrasound contrast agent with high contrast efficacy consists of a fluorinated gas bubble (for example SF₆ or a perfluorocarbon such as perfluoropropane or perfluorobutane) coated with a mono or multilayer phospholipid membrane. The particle size will generally be around 4 micrometer with very few particles larger than 10 micrometer in diameter. The main indications for such a typical product in the future may be cardiac imaging (cardiac perfusion examinations) and liver imaging.

Nuclear medicine imaging modalities are based upon administration of radioactive isotopes followed by detection of the isotopes, e.g. using gamma camera or positron emission tomography (PET). The most frequently used examination is gamma camera detection of 99-

technetium in the form of a chelate, for example a technetium phosphonate chelate for bone scintigraphy.

Light imaging methods are performed using contrast agents that absorb and/or emit light (generally near infrared light).

MSI methods may be performed without contrast agents; however, contrast agents based on magnetic materials improve this technique substantially.

APT based methods can also be performed (like for example thallium scans) without use of contrast agents; again however, contrast agents based on physiologically acceptable ions or other agents with effect on conductivity improve the diagnostic utility of APT.

All these different modalities complement each other with regard to diagnosis based on morphology/anatomy.

However, there has been a great interest in measurement and quantification of various physiological parameters. (See for example J. Magn. Reson. Imaging 1997, 7, 82-90 for a review on physiologic measurements by contrast enhanced MR imaging).

Various methods for measurements of physiologically important parameters have been described in the scientific literature: tissue pH has been measured using near infrared reflectance spectroscopy (J. Clin. Monit. 1996, 12, 387-95); intratumor pH has been measured using ^{19}F magnetic resonance spectroscopy (Invest. Radiol. 1996, 31, 680-9); 6-fluoropyridoxal polymer conjugates have been suggested as ^{19}F pH indicators for magnetic resonance spectroscopy (Bioconjug. Chem. 1996, 7, 536-40); spectral imaging microscopy has been used for simultaneous measurements of intracellular pH and Ca^{2+} in insulin-secreting cells (Am. J. Physiol. 1996, 270, 1438-46); fluorescence ratio imaging has been used for measurement of interstitial pH in solid tumors (Br. J. Cancer 1996, 74, 1206-15); a fluorinated pH probe for ^{19}F magnetic resonance spectroscopy has been used for in

vivo pH measurement after hyperthermic treatment of tumors in mice (Acta Radiol. 1996, 3, 5363-4); P-NMR has been used for analysis of intracellular free magnesium and pH in erythrocytes (J. Soc. Gynecol. Investig. 1996, 3, 66-70); intracellular pH has been estimated in developing rodent embryos using computer imaging techniques (Teratology, 1995, 52, 160-8); biscarboxyethyl carboxyfluorescein has been evaluated as in vivo fluorescent pH indicator (J. Photochem. Photobiol. B. 1995, 227, 302-8); the effect of blood flow modification on intra- and extracellular pH has been measured by ³¹P magnetic resonance spectroscopy in murine tumors (Br. J. Cancer, 1995, 72, 905-11); intracellular Ca²⁺, pH and mitochondrial function in cultures of rabbit corneal tissue have been studied by digitized fluorescence imaging (In Vitro Cell Biol. Anim. 1995, 31, 499-507); a dual-emission fluorophore has been evaluated for fluorescence spectroscopy of pH in vivo (J. Photochem. Photobiol. B. 1995, 28, 19-23); nuclear magnetic resonance spectroscopy has been used to study lactate efflux and intracellular pH during hypoxia in rat cerebral cortex (Neurosci. Lett. 1994, 178, 111-4); ³¹P NMR spectroscopy has been used for imaging of phosphoenergetic state and intracellular pH in human calf muscles after exercise (Magn. Reson. Imaging 1994, 12, 1121-6); multinuclear NMR spectroscopy has been used for studies of regulation of intracellular pH in neuronal and glial tumour cells (NMR Biomed. 1994, 7, 157-166), 5,6-carboxyfluorescein has been used as a pH sensitive fluorescent probe for in vivo pH measurement (Photochem. Photobiol. 1994, 60, 274-9); a fluorinated pH-probe has been used for non-invasive in vivo pH measurements (Invest. Radiol. 1994, 29, 220-2); fluorescence ratio imaging microscopy has been used for non-invasive measurement of interstitial pH profiles in normal and neoplastic tissue (Cancer Res. 1994, 54, 5670-4); 6-fluoro-pyridoxol has been used as probe of

cellular pH using ^1H NMR spectroscopy (FEBS Lett. 1994, 349, 234-8); lactate and pH have been mapped in calf muscles of rats during ischemia/reperfusion assessed by in vivo proton and phosphorus magnetic resonance chemical shift imaging (Invest. Radiol. 1994, 29, 217-23); nuclear magnetic resonance spectroscopy has been used for measurement of in vivo and ex vivo pH (Eur. J. Lab. Med. 1996, 4, 143-156); seminaphthofluorescein-calcein has been tested as fluorescent probe for determination of intracellular pH by simultaneous dual-emission imaging laser scanning confocal microscopy (J. Cell Physiol. 1995, 164, 9-16); ampholytic dyes have been proposed for spectroscopic determination of pH in electrofocusing (J. Chromatogr. A 1995, 695, 113-122); EPR spectroscopy has been used for direct and continuous determination of pH values in nontransparent water-in-oil systems (Eur. J. Pharm. Sci. 1995, 3, 21-6); intracellular Ca^{2+} and pH have been imaged simultaneously in glomerular epithelial cells (Am. J. Physiol. Cell Physiol. 1993, 46, 216-230); fluorinated macromolecular probes have been evaluated for non-invasive assessment of pH by magnetic resonance spectroscopy (Bioorg. Med. Chem. Lett. 1993, 2, 187-192); pH has been mapped in living tissue by application of in vivo ^{31}P NMR chemical shift imaging (Magn. Res. Med. 1993, 29, 249-251); fluorescence spectroscopy has been used to measure temperature dependent aggregation of pH-sensitive phosphatidyl ethanolamine oleic acid-cholesterol liposomes (Anal. Biochem. 1992, 207, 109-113); ^{13}C NMR spectroscopy has been used to determine intracellular pH (Am. J. Physiol. Cell Physiol. 1993, 264, C755-C760); ^{31}P NMR chemical shift imaging has been used for pH mapping of living tissue (Magn. Reson. Med. 1993, 29, 249-251); fluorescent probe and ^{31}P NMR spectroscopy have been compared for measurement of the intracellular pH of propionibacterium acnes (Can. J. Microbiol. 1993, 39, 180-6); panoramic imaging of brain pH and CBF has been

performed during penicillin and metrazole induced status epilepticus (Epilepsy Res. 1992, 13, 49-58); nuclear magnetic resonance spectroscopy has been used to study energy metabolism, intracellular pH and free Mg concentration in the brain of transgenic mice (J. Neurochem. 1992, 58, 831-6); the pH dependence of 5-fluorouracil uptake has been observed by in vivo ^1H and ^{19}F nuclear magnetic spectroscopy (Cancer Res. 1991, 51, 5770-3); ^31P magnetic resonance spectroscopy has been used to study tumor pH and response to chemotherapy in non-Hodkin's lymphoma (Br. J. Radiol. 1991, 64, 923-8); ^31P magnetic resonance spectroscopy and microelectrodes have been used to evaluate dose-dependent thermal response of tumor pH and energy metabolism (Radiat. Res. 1991, 127, 177-183); hepatic intracellular pH has been studied in vivo by ^{31}P NMR spectroscopy (Magn. Reson. Med. 1991, 19, 386-392); the relationship between vertebral intraosseous pressure, pH, pO_2 , pCO_2 and magnetic imaging signal inhomogeneity has been evaluated in a patient with back pain (Spine 1991, 16, 239-242); the effect of hypoxia on phosphorus metabolites and intracellular pH in the fetal rat brain have been studied by ^{31}P NMR spectroscopy (J. Physiol. 1990, 430, 98P); brain pH in head injury has been evaluated using image-guided ^{31}P magnetic resonance spectroscopy (Ann. Neurol. 1990, 28, 661-7); Se-labeled tertiary amines have been prepared and evaluated as brain pH imaging agents (Nucl. Med. Biol. Int. J. Radiat. Appl. Instrum. Part B 1990, 17, 601-7); ^1H , ^{31}P and ^{13}C nuclear magnetic resonance spectroscopy have been used to study cerebral energy metabolism and intracellular pH during severe hypoxia and recovery in the guinea pig cerebral cortex in vitro (J. Radiat. Appl. Instrum. Part B 1990, 26, 356-369); development of a pH-sensitive contrast agent for ^1H NMR imaging has been reported (Magn. Reson. Med. 1987, 5, 302-5); and there have been other references to ^{31}P NMR studies of pH, see for example Biomed. Res.

- 8 -

(Japan) 1989 10, Suppl. 3, 587-597, J. Cereb. Blood Flow Metab. 1990, 10, 221-6, Br. J. Radiol. 1990, 63, 120-4, Pediatr. Res. 1989, 25, 440-4, Radiology 1989, 170, 873-8, Cereb. Blood Flow Metab. 1988, 8, 816-821, J. Neuro. Chem. 1988, U51U, 1501-9 and Am. Heart J. 1988, 116 701-8. WO98/41241 of Nihon Schering discusses MRI techniques which utilise polymers in the monitoring of pH.

One important physiological parameter of great medical interest has been temperature; temperature has been measured by electron paramagnetic resonance spectroscopy (J. Biomech. Eng. 1996, 118, 193-200), an ytterbium chelate has been used as a temperature sensitive probe for MR spectroscopy (Magn. Res. Med. 1996, 35, 648-651), fast imaging techniques have been evaluated in MRI for temperature imaging (J. Magn. Reson. B, 1996, 112, 86-90), ³¹P and ¹H magnetic resonance spectroscopy has been used to study relationship between brain temperature and energy utilization rate in vivo (Pediatr. Res. 1995, 38, 919-925), local brain temperature has been estimated in vivo by ¹H NMR spectroscopy (J. Neurochem. 1995, 38, 1995, 1224-30), magnetic resonance has been used to follow temperature changes during interstitial microwave heating (Med. Phys. 1997, 24, 269-277), the temperature dependence of canine brain tissue diffusion coefficient has been measured in vivo using magnetic resonance echoplanar imaging (Int. J. Hyperthermia 1995, 11, 73-86), temperature dependent ultrasound colour flow Doppler imaging has been carried out of experimental tumours in rabbits (Ultrasound Med. Biol. 1993, 19, 221-9), electrical impedance tomography has been proposed for temperature measurement (Trans ASME J. Biochem. Eng. 1996, 118, 193-200), temperature measurement has been carried out in vivo using a temperature-sensitive lanthanide complex and ¹H magnetic resonance spectroscopy (Magn. Res. Med. 1996, 35,

364-9), body temperature imaging by impedance CT has been carried out (Med. Imag. Tech. (Japan) 1995, 13, 696-702), temperature imaging has been carried out inside the human body using microwaves (Med. Imag. Techn. (Japan) 1995, 13, 691-5), in vivo oxygen tension and temperature have been determined simultaneously using ^1F NMR spectroscopy of perfluorocarbon (Mag. Res. Med. 1993, 29, 296-302), measurement of in vivo pH in normal and tumor tissue has been carried out by localized spectroscopy using a fluorescent marker (Optical Eng. 1993, 32, 239-43), microwave temperature imaging has been proposed (IEEE Trans. Med. Imag. (USA) 1992, 4, 457-69), non-invasive temperature mapping during hyperthermia has been carried out by MR imaging of molecular diffusion (Proceedings of the Annual International Conference of the IEEE 1988, 342-343). There have been other reports of non-invasive and minimally invasive methods for the early detection of disease states by MRI, positron emission tomography, EEG imaging, MEG imaging, SPECT, electrical impedance tomography (APT), ECG imaging and optical diffusion tomography, see for example Proceedings of the SPIE - The International Society for Optical Engineering (USA) 1887 (1993).

The following, predominantly MRI based, techniques have also been reported in the measurement of temperature and temperature changes; Med. Phys 1997, 24(2), 269-277, Int. J. Hyperthermia 1995, 11(5), 409-424, Int. J. Hyperthermia 1992, 8(2), 253-262, Int. J. Hyperthermia (1994), 10(3), 389-394, Radiologe 1998, 38, 200-209, Med Phys 1997, 24(12), 1899-1906, JMRI 1998, 8, 128-135, JMRI, 1998, 8, 160-164, JMRI 1998, 8, 165-174, MRM 1995, 34, 359-367, MRM 1995, 33, 729-731, MRM 1995, 33, 74-81, Radiology 1998, 208, 789-794, JMRI 1996, 7, 226-229, JMRI 1997, 8 188-196, JMRI 1998, 8, 197-202, JMRI 1998, 8, 31-39, JMRI 1998, 8, 121-127, JMRI 1998, 8, 493-502, Int. J. Radiation Oncology Biol. Phys. 1998,

- 10 -

40(i), 815-822, Int. J. Hyperthermia 1998, 14(5), 479-493, Radiology 1995, 196, 725-733, Advances in Radiation Therapy 1998 Eds. Mittal, Purdy and Ang, Kluwer Academic Publishers, Chapter 10, pp. 213-245.

Several patents and patent applications which relate to physiological imaging have been published: use of macrocyclic metal complexes as temperature probes for the determination of body temperature using spectroscopic methods with reduced background signals (WO94/27977); new fluorine containing macrocyclic metal complexes from tetraazadodecane derivatives useful for measuring tissue temperature from NMR chemical shift values, and as contrast agents for X-ray or NMR diagnosis (WO94/27978); determining and imaging of temperature change in human body using diffusion coefficients obtained by NMR to determine absolute temperature for individual points of body and temperature differences (WO90/02321); thermographic imaging using a temperature dependent paramagnetic material in an ESR enhanced magnetic resonance imaging apparatus (WO90/02343); fluorosubstituted benzene derivatives useful as agents for in vivo NMR diagnosis, e.g. for measurement of tissue specific pH temperature, redox potentials, etc. (EP-A-368429); a magnetic resonance pulsed heat system for selectively heating a region of a subject that uses pulsed heat from focussed ultrasound equipment to destroy tumor tissue and MRI to provide fast scan images to monitor tissue and temperature with a diffusion sensitive pulse sequence (US-A-5247935); a magnetic resonance pulsed heat system for selectively heating tissue - surgery is performed using localised heating of tissue guided by and monitored by temperature sensitive magnetic resonance imaging and body tissue is heated using a magnetic resonance imaging system having a source and a probe containing a magnetic imaging coil and heating imaging rf source (US-A-5323778); apparatus for hyperthermia

treatment of cancer comprising a combined hyperthermia and MRI probe to simultaneously heat a malignant area and monitor temperature, with a filter to isolate signals (WO91/07132); and a temperature measurement method using tomographic techniques of magnetic resonance imaging to measure the temperature of a region indirectly from an intensity change of magnetic resonance signal (US-A-5207222).

The present invention however is based on the understanding that particulate contrast agents may be produced in which the matrix or membrane material for the particles is responsive to a particular physiological parameter resulting in a change in the contrast efficacy of the contrast agent which may be correlated to that physiological parameter.

Thus viewed from one aspect the invention provides a method of imaging of an animate human or non-human animal body, which method comprises: administering parenterally to said body a particulate material comprising a matrix or membrane material and at least one contrast generating species, which matrix or membrane material is responsive to a pre-selected physiological parameter whereby to alter the contrast efficacy of said species in response to a change in the value of said parameter; generating image data of at least part of said body in which said species is present; and generating therefrom a signal indicative of the value or variation of said parameter in said part of said body.

Viewed from a further aspect the invention provides a parenterally administrable contrast medium for imaging of a physiological parameter, said medium comprising a particulate material the particles whereof comprise a matrix or membrane material and at least one contrast generating species, said matrix or membrane material being responsive to said physiological parameter to cause the contrast efficacy of said contrast generating

- 10 -

species to vary in response to said parameter. In a particularly preferred embodiment, the matrix or membrane material comprises a lipid or lipid mixture having a T_c value between 35 and 80°C, preferably between 37 and 45°C, more preferably between 38 and 43°C (T_c is defined as the gel-to-liquid crystalline phase temperature). In a further preferred embodiment, the matrix or membrane material comprises peptides or one or more polymers.

Viewed from a still further aspect the invention provides the use of a contrast generating species for the manufacture of a particulate contrast medium for use in a method of diagnosis comprising generating a signal indicative of the value of said physiological parameter, the particles of said contrast medium comprising a matrix or membrane material and at least one contrast generating species, said matrix or membrane material being responsive to said physiological parameter to cause the contrast efficacy of said contrast generating species to vary in response to said parameter.

In the method of the invention, the image data generated may if desired be presented as a two or more dimensional spatial image, alternatively they may be presented as a temporal image, again in two or more dimensions. However in the extreme the data may simply provide one or more image values (e.g. numerical values) which either directly or indirectly may be used to provide quantitative or qualitative information (a signal) indicative of the value of the parameter under study. The image data may if desired be presented in visualizable form but alternatively they may simply be a set of data points which are collected and operated on to produce the signal without a visible image actually being generated. The signal indicative of the value of the parameter under study may likewise be generated in the form of a visible image, e.g. a map of the parameter value within the body, or a chart showing variation of

- 13 -

the parameter value with time, or it may simply be a calculated numerical value for the parameter or an indication that the parameter is below or above a particular threshold value. Desirably, however, the signal provides a quantitative or at least semi-quantitative value for the parameter, e.g. either in a region of interest or in a plurality of regions of interest in the body, for example providing a spatial and/or temporal map of the parameter within at least a portion of the body.

Data relating to a physiological parameter may not necessarily also contain information relating to the anatomy of the animal body and thus, a further aspect of the invention relates to the combination of traditional anatomical imaging with physiological imaging to obtain two images, one containing information about a physiological parameter and the other containing anatomical information. The two images may be combined to give one image with both anatomical and physiological information.

Thus, according to a further aspect is provided a method of imaging of an animate human or non-human animal body, which method comprises:

administering parenterally to said body at least one contrast generating species the contrast efficacy whereof is responsive to a change in value of a pre-selected physiological parameter;

generating image data of at least part of said body in which said species is present; and

generating therefrom a signal indicative of the value or variation of said parameter in said part of said body and also generating an anatomical image of the same part of the animal body.

The additional use of anatomical information may aid interpretation of the physiological data. An image generated in response to a physiological parameter, a 'physiological image', may be formed using any of the

imaging methods and or contrast media described herein. This physiological image can be combined with a conventional image obtained with or without a contrast agent. Suitable contrast agents for use with traditional anatomical imaging are well known in the art for all types of imaging techniques, MRI, X-ray, ultrasound, light and nuclear imaging etc. and many suitable contrast agents for anatomical imaging are discussed herein.

The imaging technique used to obtain physiological data may be the same or different to the imaging technique used to obtain the anatomical image. In a preferred embodiment the imaging technique will be the same, MRI being particularly suitable.

Two separate contrast agents may be used, one for physiological imaging and one for traditional imaging. The two agents can be injected sequentially and the body scanned sequentially with respect to the appropriate imaging techniques and optionally the two images which are generated are then combined. In an alternative embodiment, a single multi-functional contrast agent may be used which is capable of providing both physiological and anatomical information. A multi-functional MRI contrast agent may be used, wherein one of its functions responds to a physiological parameter while a second function provides anatomical information. Although a single contrast agent is applied, the body may be scanned twice and the resulting two images combined.

In a further alternative embodiment a multi-functional contrast agent may be used wherein the components of the agent function as contrast agents for different imaging techniques. Thus, the contrast agent may contain microbubbles to provide contrast in ultrasound imaging and paramagnetic complexes for MRI, one of these components being responsive to a physiological parameter. Again, the images obtained by scanning according to the two imaging techniques may be

combined.

By way of a further example, MRI with hyperpolarised substances will tend to provide good physiological information relating to e.g. pH, temperature or pressure but little or no anatomical information. Thus, the hyperpolarised MR image is advantageously combined with an anatomical image, e.g. by superimposing the images. The two images may be generated separately or at the same time.

The combination of physiological and anatomical imaging may be used to investigate all parts of the human or non-human animal body and any of the physiological parameters discussed herein, particularly pH and temperature. Where the physiological parameter is temperature, changes in the value of the parameter, i.e. temperature changes, may be caused by intrinsic or extrinsic means. Intrinsic means will include cancer, cardiovascular disease and inflammation while extrinsic means include hyperthermia (external heating) treatment. Thus, the physiological contrast agent may be a contrast agent for hyperthermia.

The imaging technique used in the method of the invention may be any technique capable of use in conjunction with contrast agents, e.g. X-ray (e.g. CT scanning), MRI, MRS, MR microscopy, ESR imaging, ESR spectroscopy, Mössbauer imaging, ultrasound, light imaging, nuclear imaging (e.g. scintigraphy, PET or SPECT), MSI, APT, etc. In magnetic resonance techniques, signal strength or chemical shift or both may typically be studied. Preferably, the technique used will be an X-ray, MRI, ultrasound, light imaging or nuclear imaging technique (e.g. scintigraphy), in particular an MRI or ultrasound technique. The particulate contrast agent used will accordingly be or contain a material capable of having a contrast or signal generating effect in the particular imaging modality selected, e.g. a gas or gas precursor, a

- 16 -

paramagnetic, ferromagnetic, ferrimagnetic or superparamagnetic material or a precursor therefor, hyperpolarized nmr active (i.e. non zero nuclear spin) nuclei (e.g. noble gas or ^{13}C nuclei), a radionuclide, a chromophore, (which term is used to include fluorescent and phosphorescent materials as well as light absorbers) or a precursor therefor, an ionic species, etc.

The physiological parameter studied using the method of the invention may be any physiochemical parameter capable of affecting the matrix or membrane material of the contrast agent, e.g. pressure, temperature, pH, oxygen tension, carbon dioxide tension, enzyme activity, metabolite concentration, tissue electrical activity, tissue diffusion, ion concentration, particularly Mg^{2+} , Ca^{2+} and Zn^{2+} , etc. Preferably however it will be selected from blood parameters, e.g. pressure, temperature and pH, in particular in the vasculature rather than the chambers of the heart. Where temperature is being measured, changes may be due to intrinsic factors such as disease or because of external factors, i.e. hyperthermia. It is not envisaged that the parameter be one which does not affect the membrane or matrix, for example flow rate or perfusion density.

A key part of the present invention is that the contrast agent particles should comprise a membrane or matrix material which is responsive to the physiological parameter under investigation so as to alter the contrast efficacy of the contrast agent. The manner in which the membrane or matrix responds will depend on the particular combination of imaging modality, physiological parameter and contrast generating material selected. Typically however the response might involve a change in membrane or matrix permeability to one or more species (e.g. water or gases), chemical or physical breakdown of the membrane or matrix material, generation of a contrast generating material, cleavage of

functional groups from a contrast generating material thereby changing its contrast generating ability, alteration of oxidation state in a contrast generating material thereby changing its contrast generating ability, etc. Such a response may thus for example involve release from the particulate contrast agent of water-soluble contrast generating moieties that are capable of being taken up into the extracellular fluid outside the vasculature. Particular examples of physiological parameter responsive particulate contrast agents will be described in greater detail below.

Thus one embodiment of the invention relates to thermosensitive paramagnetic particulate compositions for temperature MRI-mapping of the human body. Another embodiment of the invention relates to the use of thermosensitive particulate gas compositions as an ultrasound-based in vivo thermometer.

Yet another embodiment of the invention relates to radioactive compositions for temperature mapping in the human body. Another embodiment of the present invention relates to thermosensitive particulate compositions containing water-soluble X-ray contrast agents for mapping of temperature in the human body.

Still another aspect of the present invention relates to particulate compositions containing near infrared dyes for light imaging based temperature mapping in the body.

Another aspect of the present invention is to use one or more of the thermosensitive particulate compositions for temperature mapping in imaging guided hyperthermia treatment.

Another embodiment of the present invention relates to pH sensitive particulate compositions for determination of pH in the body. By way of example the active contrast agent (or indicator or probe) may be a paramagnetic, magnetic or fluorinated compound detectable by MRI. The active contrast agent (or

- 18 -

indicator or probe) may be a gas or a gas generating substance for detection by ultrasound, it may be a radioactive substance for detection by scintigraphy, SPECT or PET, or it may be a fluorescent dye, a near infrared dye, a UV dye or another dye that can be detected in vivo in light imaging or light detection methods.

Yet another embodiment of the invention relates to particulate compositions as contrast agents or as in vivo indicators or probes for detection of oxygen concentration/tension in the tissue using modalities such as ultrasound, MRI, Overhauser MRI and ESR.

Another embodiment of the present invention relates to particulate compositions as contrast agents or as in vivo indicators or probes for detecting pressure, turbulence, viscosity, enzyme activity, ion concentrations, metabolite diffusion coefficients, elasticity and flexibility.

Another aspect of the present invention relates to particulate compositions as contrast agents or as in vivo indicators or probes in combination with a targeting ligand, wherein said targeting ligand targets cells or receptors selected from the group consisting of myocardial cells, endothelial cells, epithelial cells, tumor cells, brain cells, and the glycoprotein GPIIb/IIIa receptor, for detection of changes in physiological parameters and/or quantification/semiquantification of physiological parameters relevant for diagnosis of disease.

Further examples of targeting ligands which can be used are:

i) Antibodies, which can be used as vectors for a very wide range of targets, and which have advantageous properties such as very high specificity, high affinity (if desired), the possibility of modifying affinity according to need etc. Whether or not antibodies will be bioactive will depend on the specific vector/target

combination. Both conventional and genetically engineered antibodies may be employed, the latter permitting engineering of antibodies to particular needs, e.g. as regards affinity and specificity. The use of human antibodies may be preferred to avoid possible immune reactions against the vector molecule.

A further useful class of antibodies comprises so-called bispecific antibodies, i.e. antibodies having specificity for two different target molecules in one antibody molecule. Such antibodies may, for example, be useful in promoting formation of bubble clusters and may also be used for various therapeutic purposes, e.g. for carrying toxic moieties to the target. Various aspects of bispecific antibodies are described by McGuinness, B.T. et al. in *Nat. Biotechnol.* (1996) **14**, 1149-1154; by George, A.J. et al. in *J. Immunol.* (1994) **152**, 1802-1811; by Bonardi et al. in *Cancer Res.* (1993) **53**, 3015-3021; and by French, R.R. et al. in *Cancer Res.* (1991) **51**, 2353-2361.

ii) Cell adhesion molecules, their receptors, cytokines, growth factors, peptide hormones and pieces thereof. Such vectors/targeting ligands rely on normal biological protein-protein interactions with target molecule receptors, and so in many cases will generate a biological response on binding with the targets and thus be bioactive; this may be a relatively insignificant concern with vectors which target proteoglycans.

iii) Non-peptide agonists/antagonists or non-bioactive binders of receptors for cell adhesion molecules, cytokines, growth factors and peptide hormones. This category may include non-bioactive vectors which will be neither agonists nor antagonist but which may nonetheless exhibit valuable targeting ability.

iv) Oligonucleotides and modified oligonucleotides which bind DNA or RNA through Watson-Crick or other types of base-pairing. DNA is usually only present in

- 20 -

extracellular space as a consequence of cell damage, so that such oligonucleotides, which will usually be non-bioactive, may be useful in, for example, targeting of necrotic regions, which are associated with many different pathological conditions. Oligonucleotides may also be designed to bind to specific DNA- or RNA-binding proteins, for example transcription factors which are very often highly overexpressed or activated in tumour cells or in activated immune or endothelial cells. Combinatorial libraries may be used to select oligonucleotides which bind specifically to possible target molecules (from proteins to caffeine) and which therefore may be employed as vectors for targeting.

v) DNA-binding drugs may behave similarly to oligonucleotides, but may exhibit biological activity and/or toxic effects if taken up by cells.

vi) Various small molecules, including bioactive compounds known to bind to biological receptors of various kinds. Such vectors or their targets may be used to generate non-bioactive compounds binding to the same targets.

vii) Targeting ligands may be selected from combinatorial libraries without necessarily knowing the exact molecular target, by functionally selecting (in vitro, ex vivo or in vivo) for molecules binding to the region/structure to be imaged.

ix) Proteins or peptides which bind to glucosamino-glycan side chains e.g. heparan sulphate, including glucosaminoglycan-binding portions of larger molecules, since binding to such glucosaminoglycans side chains does not result in a biological response. Proteoglycans are not found on red blood cells, thus eliminating undesirable adsorption to these cells.

The particulate contrast agent may thus be used for quantification/semi-quantification of a physiological parameter which is relevant for diagnosis of disease. The particulate contrast agent may be triggered into

- 21 -

giving a measurable signal difference either by the target parameter itself (e.g. the local temperature, pH or pressure or by binding to the particular cell surface receptors of interest) or by a chemical or biological response of the target parameter (e.g. release of enzymes or local variation in pH or temperature due to cellular reactions). The particulate agent may thus respond to, identify and/or quantitatively or semi-quantitatively determine bacteria, viruses, antibodies, enzymes, drugs, toxins, etc.

Another aspect of the present invention relates to intravenous particulate compositions as contrast agents or as in vivo indicators or probes with long vascular half life (reduced liver uptake) for detection of changes in physiological parameters and/or quantification/semiquantification of physiological parameters relevant for diagnosis of disease.

The particulate contrast agent used according to the invention may be a solid material, a porous material, a liquid crystal material, a gel, a plastic material, a material having one or more walls or membranes or liquid particles, e.g. emulsion droplets or gas based particles, e.g. micro bubbles. The particles can also be thermodynamically stabilised, e.g. micro emulsion droplets or surfactant micelles. The chemical composition of the particulate material can be one simple chemical compound or a mixture of two or more chemical compounds. Generally it will comprise two or more different chemical entities, at least one of which is a matrix or membrane forming material and at least one other of which is a contrast generating species. The composition can consist of solid material(s) only or it may be a mixture of different solids/liquids/gases. The particulate will generally have a mean particle size (e.g. as determined by particle size analyzers such as laser light scattering apparatus or Coulter counters) in the range 0.001 to 20 μ m, more preferably 0.01 to 10 μ m,

especially 0.05 to 7 μm . Such particles are often described in the literature as particles, colloids, emulsions, droplets, microcrystals, nanocrystals, microparticles, nanoparticles, vesicles, liposomes, bubbles, microspheres, microbubbles, coated particles, microballons and the like.

The term "polymer" as used herein refers to any chemical compound with more than 10 repeating units. A polymer can be naturally occurring, synthetic, or semisynthetic. Semisynthetic polymers are polymers that are produced by synthetic modification of naturally occurring polymers. Compounds with 2 to 10 repeating units are herein generally referred to as "oligomers" and likewise may be natural, synthetic or semisynthetic.

The term "surface active compound" or "surfactant" is used herein to refer to any chemical compound having at least one hydrophilic functional group and at least one hydrophobic (lipophilic) group. In a multiphase system, surface active compounds will commonly accumulate at the interface.

The term "lipid" is used herein to refer to naturally-occurring compounds, synthetic compounds and semisynthetic compounds which are surface active compounds and have structures similar to fatty acids, waxes, mono-, di- or tri-glycerides, glycolipids, phospholipids, higher (C_{10} or greater) aliphatic alcohols, terpenes and steroids.

The term "gas" is used herein to refer to any compound or a mixture of compounds with sufficiently high vapor pressure to be at least partly in the gas phase at 37°C.

When the imaging modality used according to the invention is ultrasound, the contrast generating species in the contrast agent will preferably consist of one or more encapsulated gases and/or one or more encapsulated gas precursors. This contrast generating species is able to interact with the surroundings so that the

contrast agent gives information about one or more physiological parameters generally as a result of an interaction between the surroundings and the encapsulation material, if necessary followed by changes related to the gas/gas-precursor. However gaseous contrast generating species may be used in other imaging modalities, such as MRI and X-ray for example.

Typical examples of gas types that change contrast property as a result of the physiological parameters in the surrounding tissue include: gases that are generated from a precursor as a result for example of pH, temperature or pressure changes, e.g. as a result of a chemical reaction, as a result of the boiling point of the gas, or as a result of a change of solubility; gases that compete with blood gases for absorption or adsorption sites within the matrix or membrane material; gases that change properties (e.g. lose hyperpolarization or change other magnetic properties) upon contact with body fluids or components, including dissolved components, thereof; gas molecules sensitive to pH; gases that change properties/volume with temperature; gases that change volume as a result of surrounding gas (e.g. oxygen tension); etc.

Preferred gases include hydrogen, oxygen, nitrogen, noble gases (including hyperpolarized gases), carbon dioxide, fluorinated gases (e.g. sulphur hexafluoride, fluorohydrocarbons, perfluorocarbons and other fluorinated halogenated organic compounds in gas phase), and low molecular weight hydrocarbons. Preferred gases also include any pharmaceutically acceptable gas mixture like for example air and air/perfluorocarbon mixtures. Preferably, the perfluorocarbon gas is selected from perfluoromethane, perfluoroethane, perfluoropropanes and perfluorobutanes. Any physiologically acceptable gas precursor can be used. Among suitable gas precursors are compounds that form a gas as a result of a chemical reaction (for example compounds sensitive to pH, for

example carbonic acid, aminomalononic acid or other acceptable pH sensitive gas generating substances). Other suitable gas precursors are compounds that form a gas as a result of other physiological conditions like for example temperature, oxygen, enzymes or other physiological parameters/compounds relevant for body tissue (whether in the normal or diseased state) or which are activated to a gas forming state as a result of an interaction with an external stimulus (e.g. photo-activation, sono-activation etc.).

The encapsulation material can be any material such as for example lipids, phospholipids, surfactants, proteins, oligomers and polymers. Such materials may be chosen to dissolve, melt, collapse, weaken, increase porosity, or otherwise break down, change phase or change size (e.g. by aggregation due to change in surface charge, for example in response to local Ca^{2+} and/or Mg^{2+} concentration) in response to the physiological parameter, e.g. to allow release of the contrast generating species into the surrounding fluid, or to allow body fluid or components thereof to come into contact with the contrast generating species, or to raise contrast agent species local concentration above the detection limit, etc. In this way the contrast generating effect of the contrast generating species may be dispersed (e.g. into the extracellular fluid space), switched on or increased (e.g. by generation of a contrast generating species such as a gas or by increasing water contact (for a positive (T_1 effect) MR contrast agent such as a gadolinium chelate)), or switched off or decreased (e.g. by destruction of the compartmentalization required for a negative (T_2 effect) MR contrast agent such as a dysprosium chelate, or by quenching of a radical or depolarization of a hyperpolarized nucleus or dissolution of a blood soluble gas). Moreover a porous solid matrix, e.g. a zeolite, may be impregnated with the contrast generating species

with the pore mouths then being closed off totally or partially using a material which breaks down, melts or dissolves when the relevant physiological parameter (e.g. pH, temperature, enzyme concentration) in the surrounding body fluid is above or below a pre-set value.

The particulate contrast agent used according to the invention may respond to physiological parameters in several different ways. In one aspect, the particulate contrast agent may respond to physiological parameters by accumulation in the area where a certain value for a particular parameter is fulfilled, compared to areas where it is not. In another aspect of the invention, the particulate contrast agent responds by accumulation in areas where the physiological parameter value is not fulfilled. In yet another aspect of the invention, the particulate contrast agent responds to a given parameter by disintegration, the disintegration being dissolution or chemical breakdown. Especially advantageous is a response to a physiological parameter by leakage or other transport means in/out of the particles. The opposite situation where the response to a physiological parameter is to prevent dissolution/leakage by attaining an increase in stability/reduction in membrane transport compared to particles in areas where a threshold value for a given parameter is not fulfilled, is also a preferred aspect of the present invention. This type of response is advantageous since a time course may lead to a reduction in contrast by elimination from the organ in areas where the threshold value for the parameter is not fulfilled, while the contrast remains in the area of interest.

When a particulate composition responds by disintegration or transport, changes in contrast effect may be achieved by exposing otherwise invisible/shielded contrast agents, altering the distribution of contrast agents or, when the contrast agent is the particle

- 26 -

itself (as in ultrasound contrast agents), destroying the contrast giving property. Especially advantageous are particulate compositions where the contrast effect is gained by interaction with the environment. In this case, both transport of the contrast agent and transport of the actual environmental component may be utilized for detection of physiological parameters. An example is MRI contrast agents where an increased degree of water access/transport to the contrast agent leads to the measured contrast enhancement. In this case, response to a physiological parameter may be an increased rate of water transport in/out of the particulate.

The leakage or an increased transport rate of solutes in/out of a particulate may be accomplished in a variety of ways. All kinds of phase transitions may be utilized to induce leakage/transport. For instance, a solid particle/membrane may become leaky when it is melted, the process being sensitive to temperature. Phase transitions involving a gas phase may be used to respond to pressure as a physiological parameter. An especially useful aspect of the present invention is particles comprising liquid crystalline material as for example liposomes, niosomes or other vesicles. Liquid crystalline materials may undergo several different phase changes which may induce leakage and/or increase the transport rate of solutes or even breakdown of the particle. For example, the gel to liquid crystalline phase transition of phospholipids may increase the liposome permeability and increase the transport rate or induce leakage of solutes on heating and hence temperature sensitivity. The lamellar to reversed hexagonal phase transition will also induce leakage since the liposomes require lipids in lamellar, gel or other layered phase structure. The lamellar to reversed hexagonal phase transition may be induced by pH, electrolytes, and changes in the chemical environment

- 27 -

such as targeting, enzymes, antibodies etc. The suitable parameter to respond to may be tuned by selection of the membrane composition and processing. Other phase transitions such as lamellar to cubic phases, lamellar to microemulsion phases or lamellar to normal hexagonal phase may also be used to introduce leakage.

Gel based particles or gel-surrounding particles (e.g. particles made by coacervation) may respond to a physiological parameter by, for example, a lowering of the viscosity of the gel. Such viscosity lowering may for example be obtained by temperature, pH or electrolytes such as Ca or Mg and the particles are thus sensitive to these parameters. Such parameters may also induce phase separation in the gel particles, leading to leakage of liquid and phase separation of the polymer which comprises the gel. These mechanisms may in turn influence a parameter such as water leakage and exposure of, e.g. paramagnetic chelates to water and hence lead to a change in MRI contrast.

Particles or membranes composed of solid polymer may also respond to physiological parameters. For instance temperature may change the glass transition temperature of the polymer, and hence induce phase transitions in the polymer membrane, which in turn may influence a parameter such as water transport which influences the contrast efficacy of the contrast agent.

Particles which at least in part are composed of or stabilised by water soluble polymers e.g. peptides, may respond to physiological parameters by alteration in the peptide conformation. For instance peptides may undergo an α - helix to β - sheet transition or vice versa and hence influence a parameter which in turn effects contrast. Also transitions to/from α - helix or β - sheet to random coil may influence a parameter such as membrane permeability, particle stability against aggregation/flocculation or even fusion, or particle

- 28 -

dissolution or precipitation which in turn alters the contrast efficacy of the contrast agent.

Also of use as contrast agents when the imaging modality is ultrasounds are temperature and pressure sensitive emulsions and fluids.

Leakage may also be controlled by entities forming channels or other transport routes through the membrane of a particle. These channels may control the transport of molecules in/out of the particle, and be quite selective for, e.g., ions. For instance the protein tubulin which forms microtubules in absence of Ca^{++} may induce a higher leakage in presence of Ca^{++} than in absence of Ca^{++} and hence be Ca^{++} sensitive. Other proteins/enzymes which may control the transport of substance in/out of a vesicle, include erythrocyte anion transporter, erythrocyte glucose transporter, $\text{Na}^{+}\text{-K}^{+}$ ATPase ($\text{Na}^{+}\text{/K}^{+}$ pump), Ca^{++} - ATPase (Ca^{++} pump) and Bacteriorhodopsin (H^{+} - pump). Also biosurfactants such as iturins, esperine, bacillomycins, mycosubtilin, surfactin and similar substances may be used as membrane components to induce/prevent leakage by response to external parameters since these molecules may respond by changes in secondary and tertiary structure as well as self-assembly properties on influence from extrinsic parameters.

The contrast generating species in MR contrast agents used according to the invention will generally be a paramagnetic, superparamagnetic, ferrimagnetic or ferromagnetic compound and/or a compound containing other non zero spin nuclei than hydrogen, e.g. ^{19}F , ^{13}C , ^{15}N , ^{29}Si , ^{31}P and certain noble gases, such as ^{129}Xe or ^3He .

Preferred as paramagnetic compounds are stable free radicals, and compounds (especially chelates) of transition metal or lanthanide metals, e.g. manganese compounds, gadolinium chelates, ytterbium chelates and dysprosium chelates. Preferred magnetic (e.g.

- 29 -

superparamagnetic) compounds are γ -Fe₂O₃, Fe₃C and other iron/metal oxides with high magnetic susceptibility. Preferred fluorinated compounds are compounds with relative short F T₁-relaxation times. Other preferred fluorinated compounds according to the present invention are fluorinated pH-probes, such as compounds described in EP-0447013 of Schering A.G. and ZK-150471 described by Y. Aoki in Invest. Radiol 1996, 34, 680-689. Examples of MR contrast effective materials are well known from the patent literature, see for example the patent publications of Nycomed, Salutar, Sterling Winthrop, Schering, Squibb, Mallinckrodt, Guerbet and Bracco.

In general, there are two types of contrast generating species useful in MR contrast agents for use according to the invention: species that change contrast property as a result of the physiological parameters in the surrounding tissue; and species that are inert to physiology but change contrast properties as a result of an interaction between coating material/encapsulation material and physiology. Typical examples here will be GdDTPA, GdDTPA-BMA, GdDOTA, GdHPDO3A, PrDO3A-derivatives and Tm chelates in thermosensitive liposomes or in pH-sensitive vesicles.

Typical examples of species that change contrast property as a result of the physiological parameters in the surrounding tissue include: paramagnetic chelates that change relaxation properties and/or change chemical shift as a result of temperature, paramagnetic chelates that change coordination number and thereby relaxation properties and/or shift properties as a function of pH, paramagnetic compounds, for example manganese compounds (Mn(2+)/Mn(3+)), europium compounds (Eu(2+), Eu(3+)) and free radicals (radical, no radical) that change relaxation properties and/or shift properties as a result of oxygen tension/concentration or as a result of redox potential in the surrounding tissue, paramagnetic

and magnetic compounds that change relaxation/shift properties as a result of enzymic activity (for example with enzymatic cleavage of paramagnetic chelates from macromolecules conjugated thereto causing a change in correlation time and/or water coordination) and paramagnetic chelates that change properties as a result of concentration of ions in the tissue, e.g. due to changes in water coordination.

Paramagnetic compounds have, according to the present invention, either an effect on the relaxation times (T_1 or T_2) or an effect on chemical shift. Typical compounds that change relaxation times are gadolinium chelates, manganese compounds and superparamagnetic iron oxides. Europium chelates, on the other hand, are well-known chemical shift compounds. The effect on chemical shift is related to temperature. Based on this, macrocyclic paramagnetic chelates like 2-methoxyethyl substituted PrDO3A and 1,4,7,10-tetraazacyclododecane-1,4,7,10-tetrakis(methylene phosphonate) thulium complex have been suggested as temperature probes (see WO 94/27977 (Platzek, Schering) and C.S. Zuo et al. in J. Magn. Res. 133 53-60 (1998)). All these paramagnetic compounds can be used according to the present invention.

The contrast generating species in X-ray contrast agents for use according to the invention will generally be a gas or gas generator or a water-soluble compound containing heavy atoms (e.g. atomic number of 37 or greater), e.g. metal chelates, metal clusters, metal cluster chelates and iodinated compounds. Preferred contrast generating species include ionic and non-ionic iodinated organic aromatic compounds, in particular triiodophenyl compounds. Most preferred are approved iodine based contrast agents such as salts, e.g. sodium or meglumine salts, of iodamide, iothalamate, diatrizoate, ioxaglate and metrizoate, and non-ionics such as metrizamide (see DE-A-2031724), iopamidol (see

31 -

BE-A-836355), iohexol (see GB-A-1548594), iotrolan (see EP-A-33426), iodecimol (see EP-A-49745), iodoxanol (see EP-A-108638), ioglucol (see US-A-4314055), ioglucomide (see BE-A-846657), ioglunide (see DE-A-2456685), iogulamide (see BE-A-882309), iopromide (see DE-A-2909439), iosacol (see DE-A-3407473), iosimide (see DE-A-3001292), iotasul (see EP-A-22056), ioversol (see EP-A-83964) and ioxilan (see W087/00757).

Such contrast generating species may be incorporated into matrices or coatings that are sensitive to one or more physiological parameter.

The contrast generating species in nuclear medicine contrast agents for use according to the invention may be any radioactive compound of the type in diagnostic nuclear medicine, for example known compounds useful for scintigraphy, SPECT and PET. Typical compounds include radiolodinated compounds, ¹¹¹Indium labelled materials and ^{99m}Tc labelled compounds (for example ^{99m}TcDTPA, ^{99m}TcHIDA and ^{99m}Tc labelled polyphosphonates) and ⁵¹CrEDTA.

Such contrast generating species may be incorporated into matrices or coatings that are sensitive to one or more physiological parameter.

Contrast agents can be prepared for other imaging modalities such as light imaging, Overhauser MRI, oxygen imaging, magnetic source imaging and applied potential tomography, by encapsulation of the contrast generating species, e.g. a chromophore or fluorophore (preferably having an absorption or emission maximum in the range 600 to 1300nm, especially 700 to 1200nm), a stable free radical, a superparamagnetic particle or an ionic (preferably polyionic) species, for the respective modality into a physiologically sensitive matrix or coating.

In vivo temperature measurements have been of great interest because temperature is an important physiological parameter related to several indications including cancer, cardiovascular diseases and

inflammation. Local monitoring of temperature will also be of great value during hyperthermia treatment.

Contrast generating species can be released from the matrix/encapsulation material as a result of increased temperature and thereby change their contrast property or distribute to other tissues than the particulate product. Alternatively for an MR active temperature sensitive agent, a change in contrast efficacy may occur due to an increased permeability of the matrix/encapsulation material, and, hence, to an increased rate of water transport across the matrix/encapsulation material, even if the agent itself does not leave the matrix/encapsulation material.

Typical examples of temperature sensitive particulate materials are temperature sensitive liposomes, these being especially suitable for use with MRI. These liposomes take advantage of the fact that the membrane permeability is markedly increased at the gel-to-liquid crystal phase transition temperature (T_c) of their membrane lipids. Also, possibly depending upon the membrane properties and the nature of the MR active agent, leakage of the agent may occur. Liposomes made from specific phospholipids or a specific blend of phospholipids may be stable up to 37°C but exhibit an increased water permeability or/and leak as they pass through an area of the body in which the temperature is raised, e.g. to 40 to 45°C, as a result of a disease process or an external heating. Table 1 below shows the transition temperature of various saturated phosphatidylcholines.

Table 1

Phosphatidylcholines (PC)	Transition temperature T_c (°C)
---------------------------	-----------------------------------

- 33 -

12:0	1
13:0	14
14:0	23
15:0	33
16:0	41
17:0	48
18:0	55
19:0	60
20:0	66
21:0	72
22:0	75
23:0	79
24:0	80

Table 2 below shows the phase transition of various unsaturated phosphatidylcholines.

Table 2

Phosphatidylcholines (PC)	Transition temperature Tc (°C)
12:1	-36
18:1c9	-20
18:1t9	12
18:1c6	1
18:2	-53
18:3	60
18:4	-70

Table 3 below shows the phase transition temperature of various asymmetric phosphatidylcholines.

Table 3

Phosphatidylcholines (PC)	Transition temperature Tc (°C)
14:0-16:0	35
14:0-18:0	40
16:0-14:0	27
16:0-18:0	49
16:0-18:1	-2
16:0-22:6	-27
16:0-14:0	30
18:0-16:0	44
18:0-18:1	6
18:1-16:0	-9
18:1-18:0	9

Table 4 below shows the phase transition temperature for various saturated symmetric phosphatidylglycerols (PG) in the form of their sodium salts.

Table 4

Phosphatidylglycerols (PG)	Transition temperature T _c (°C)
12:0	-3
14:0	23
16:0	41
18:0	55

Tables 1-4 are based on information from the product catalogue of Avanti Polar Lipid Inc., USA.

Accordingly, phospholipids or blends of phospholipids may be selected to give products with the correct T_c for thermosensitive liposomes for diagnostic use. Typical blends for preparation of thermosensitive liposomes for diagnostic use are mixtures of dipalmitoylphosphatidylcholine (DPPC) and dipalmitoylphosphatidyl glycerol (DPPG) and distearylphosphatidylcholine (DSPC).

Particulate contrast agents may also respond to temperature by utilizing the conformational temperature sensitivity of certain polymer systems. An example is poly(N-isopropyl acrylamide) which phase separates at 37°C. Hence particles comprising contrast agents will become leaky dependent on temperature (see Hoffmann et al. Macromol. Symp. 118: 553-563 (1997)).

Other examples of temperature sensitive matrices/coatings are lipid suspensions/emulsions containing the contrast generating species or other particulate or particulate like formulations that release the contrast generating species or change properties as a result of changes in temperature.

If the parameter under study is capable of manipulation, e.g. by treatment with drugs, external application of heat etc., it may be used to study the efficacy of such treatment or localized such treatment may be used to cause a change in contrast efficacy which

- 36 -

in turn may be used to measure parameters such as organ perfusion. Thus for example external application of heat at, near or upstream of an organ of interest may be used to cause release from the particles of a contrast agent which may diffuse into the organ and so to detect blood perfusion (or lack of perfusion) in that organ. In this context one might administer a thermally sensitive particulate agent in connection with an external heating to follow the heat transport in parts of the body. Heat transport *in vivo* is directly connected to blood flow through the bioheat equation (J. Appl. Physiol. vol. 1, (1948), 93-122)

$$\frac{\partial T}{\partial t} r_i C_i + w_b c_b (T - T_a) = k \nabla^2 T + (Q_p + Q_m)$$

where r (kg/m^3) is the density of tissue, C_i ($\text{J/kg}^\circ\text{C}$) is the specific heat of tissue, t (s) is the time, T ($^\circ\text{C}$) is the temperature, w_b ($\text{kg/m}^3\text{s}$) is the blood perfusion, c_b ($\text{J/kg } ^\circ\text{C}$) is the specific heat of blood, T_a ($^\circ\text{C}$) is the arterial temperature, k ($\text{W/m } ^\circ\text{C}$) is the thermal conductivity of tissue, Q_p (W/m^3) is the power deposition and Q_m (W/m^3) is the local metabolic rate. Hence, the thermosensitive particulate compositions may, after a controlled, localized external heating, give a measure of blood perfusion in an organ.

The temperature response of thermosensitive MR-liposomes can in general be divided into three distinct regions:

- 'low relaxivity' region; $r_1 = r_1^{\text{low}}$; $T < T_i$; where r_1^{low} is a constant with temperature (T);
 - 'temperature active' region' $r_1(T) = f(T)$, $T_i < T < T_c$, and
 - 'high relaxivity' region; $r_1 = r_1^{\text{high}}$; $T > T_c$;
- where r_1^{high} is a constant with T (ideally, $r_1^{\text{high}} \gg r_1^{\text{low}}$).

It is possible to quantify the local temperature in the temperature active region of the liposomes, provided

three criteria are met:

1. A well defined relationship exists around T between liposomal relaxivity and temperature; i.e. $r(T) = f(T)$; $T < T < T$; where T, T is a clinically relevant temperature range. Ideally r should be a linear function of T over the range T, T .
2. The temperature active region covers a large enough temperature range.
3. The Gd concentration in tissue $[Gd]$ is known.

If $[Gd]$ is not known, even a qualitative assessment of temperature changes may prove difficult, since regions with different $[Gd]$ would have a different degree of enhancement, even if the temperature were the same. Furthermore, it would be impossible to say whether lack of enhancement after heating was due to a low local temperature or the absence of the liposomes in that region.

However the local $[Gd]$ *in vivo* can be estimated, based on the relaxation effects of the liposomes in the 'low relaxivity' state, by the following method:

1. Acquire quantitative R_1 and/or R_2/R_2^* images ($R_{1,2} = 1/T_{1,2}$) before contrast administration and after contrast administration but before hyperthermia is initiated. R_1 and R_2/R_2^* images can be routinely acquired on most state-of-the-art clinical MR systems.
2. Measure the fractional change in R_1 and/or R_2/R_2^* , $\Delta R_1 = R_1^{\text{post}} - R_1^{\text{pre}}$ in the region of interest.
3. The local Gd concentration is then given by: $[Gd] = \Delta R_1 / r_1$.
4. If r_1 is not known, the ratio of the Gd concentration between two regions is given by: $[Gd] / [Gd] = R_1 / \Delta R_1$. Alternatively, R or R^* images can be used to obtain the same Gd ratio.

In general therefore, the absolute $[Gd]$ can not be determined, unless the liposomal relaxivity in the tissue is known. However this may be fairly well

- 38 -

approximated for the r relaxivity, but not the r^* relaxivity, since this depends on tissue geometry. Nonetheless, the $[Gd]$ in one region relative to another can be estimated as described above. The relative $[Gd]$ is valuable information that can be used to adjust the signal enhancement in the image so that it reflects actual temperature changes. In order for this to be possible, one need to assume that a 'core region' exists where the temperature is above T_c . It is likely in a clinical situation that such a core region exists where the heating is most efficient surrounded by a 'penumbra' where heating is less efficient and the temperature distribution is less well defined. Now, by knowing the relative $[Gd]$ in the core versus the penumbra, the image intensity can be adjusted to compensate for any difference in $[Gd]$ in the two regions.

It is possible to estimate $[Gd_1]/[Gd_2]$ using a strongly T_1 -weighted sequence in which case change in signal intensity is almost linearly related to change in R_1 and hence $[Gd]$. This requires $TR \ll T_1$ of the target tissue. Similarly, strongly T_2 or T_2^* -weighted sequences can be used to estimate $[Gd_1]/[Gd_2]$.

Thus, when a Gd compound is encapsulated in liposomes, the resulting relaxivity (r_1, r_2) is small due to restricted water access to the paramagnetic centre. However, given that a very T_1 -sensitive sequence is used, it is still possible to detect a change in T_1 due to the presence of the liposomes prior to heating (i.e. at temperatures well below T_c). Consequently, by acquiring quantitative R_1 - or T_1 -maps of the area of interest before and after contrast injection (prior to hyperthermia treatment), the change in R_1 or T_1 enables the local Gd concentration $[Gd]$ to be determined. After heating, regional variations in $[Gd]$ can thus be accounted for; variations in contrast enhancement due to temperature differences can therefore be distinguished from variations in contrast enhancement due to

- 39 -

concentration variations.

The longitudinal relaxation rate R after contrast administration is given by:

$$R = R_0 + [Gd] \cdot r_1;$$

Where R_0 is the relaxation rate prior to contrast administration. The change in R , due to the contrast agent is therefore:

$$\Delta R = R - R_0 = [Gd] \cdot r_1;$$

After hyperthermia, a new R - or T - map is generated.

In conclusion, therefore, given that the T_1 effect of the liposomes is detectable below T_c , it is possible to map the local Gd concentration and consequently compensate for differences in contrast enhancement after heating due to local variations in Gd concentration. The R_1 or R_2 effect of the liposomes can also be used for this purpose.

In vivo pH measurements have been of great interest because pH is an important physiological parameter associated to several severe diseases. The pH value is usually reduced during cancer diseases, cardiovascular diseases like for example stroke, osteoporosis, inflammations and autoimmune diseases.

One type of pH sensitive encapsulation for diagnostic agents involves the use of pH sensitive liposomes. The general strategy is to employ pH-sensitive groups in the liposomal membrane. Such typical groups have pKa values between 4 and 5.5. Phospholipids useful for preparation of pH-sensitive diagnostic agents include diheptadecanoyl phosphatidylcholine (DHPC) in admixture with DPPC and N-palmitoyl homocystein (PHC) in different ratios (see Eur. J. Pharm. Biopharm. 1993, **22**, 97-101 for a general review on temperature and pH-sensitive liposomes).

Another type of pH-sensitive encapsulation of contrast generating species involves the use of pH-sensitive surfactants like for example N-dodecyl-2-

- 40 -

imidazole propionate (DIP) which has pKa of 6.8 (see for example Pharm. Res. 1993, 13, 404). This means that DIP at pH 7.3-7.4 (physiological) is in the non-ionized (non/low surfactant activity) form (80%) while at for example lysosomal pH (5.2) over 97% will be in the charged form.

Another means of pH-sensitive encapsulation of contrast generating species involves the use of matrix materials and/or coating materials with pKa values in the range of 4.5-7.0 so that the material is soluble or partly soluble in the charged form and insoluble or partly insoluble in the non-charged form. Such compounds can be physiologically acceptable low molecular weight compounds or physiologically acceptable polymers.

Still another means of pH-sensitive encapsulation involves the use of compounds that are chemically cleaved as a result of pH, for example polyorthoesters or polyacetals/ketals which are cleaved under acidic conditions.

Liposomes comprising phosphatidyl ethanolamines (PE) as the central component are another example of liposomes which can undergo a phase transition and become leaky when pH is reduced. pH sensitive liposomes can also be achieved by incorporation of fatty acids into phospholipid membranes.

In principle any charged particulate system where the charge is pH dependent and influences the packing of the membrane material can be used.

Access to oxygen is critical for all types of cells, and diagnostic agents for determination of oxygen concentration/tension in tissue will be of great importance in diagnosis of diseases like cancer, cardiovascular diseases, autoimmune diseases and several diseases in the central nervous system.

One type of oxygen or redox sensitive encapsulation/coating material is a material that has

- 41 -

different solubility/diffusion properties dependent on the oxygen level or the redox status; for example compounds containing a nitro-group that is reduced in vivo to an amino-group which improves solubilization of the material in reductive/low oxygen surroundings.

Determination of concentration of physiologically important ions in tissue is important for several diseases.

Types of ion concentration sensitive encapsulation materials that may be used in this regard include phospholipids, surfactants and other ion chelating materials. Negatively charged liposomes will for example bind Ca^{2+} and the membrane will change its diffusion properties and become more stiff.

An example of $\text{Ca}^{2+}/\text{Mg}^{2+}$ sensitive particulate compositions are liposomes enriched with the dimeric phospholipid cardiolipin. A cardiolipin containing membrane may undergo a lamellar to reversed hexagonal phase transition upon addition of the divalent cations since these ions bind to the cardiolipin di-phosphatidyl group.

Ca^{2+} or Mg^{2+} sensitivity may be obtained by using charge stabilised particles, e.g. solid particles, liquid particles e.g. emulsion droplets, gas particles e.g. microbubble dispersions or liposomes. Ca^{2+} or Mg^{2+} may thus induce aggregation or flocculation among the particles and by this means alter contrast effect. Ca^{2+} or Mg^{2+} sensitivity may also be obtained by using stabilising moieties for the particles which are chemically or physically influenced by Ca^{2+} or Mg^{2+} , for instance using surfactants which form water insoluble species when exposed to Ca^{2+} or Mg^{2+} and thus precipitate.

Some particles or stabilising membranes surrounding particles may also respond with a phase transition when exposed to Ca^{2+} or Mg^{2+} . An example are liquid crystalline based particles e.g. liposomes, which may respond by a lamellar to reversed hexagonal phase

transition upon addition of Ca or Mg . Also gel particles may respond easily to Ca or Mg by a significant lowering of viscosity or even phase separation of the polymer which forms basis for the gel on exposure to Ca or Mg . This viscosity reduction or phase separation may induce a change in contrast effect.

Types of enzyme sensitive encapsulation material include matrices or coatings that are degraded by enzymes, for example simple esters of low molecular weight compounds or polyesters like polyacetic acid and others.

Various metabolites may also change the properties of coating materials.

Particulates can be made sensitive to for example antibodies based on enhanced leakage due to a phase transition induced by the chemical binding between membrane molecules and the antibody. As an example, liposomes comprising N-(dinitrophenylamino- ϵ -caproyl)-phosphatidyl ethanolamine (DNP-cap-PE) become leaky due to a lamellar to reversed hexagonal phase transition when binding to anti-DNP. Another example includes liposomes comprising human glycophorin A in dioleoyl phosphatidyl ethanolamine membranes. These liposomes become leaky when immobilized antibodies are added.

A further aspect of the present invention is to use one of the above described particulate diagnostic agents together with another compound that has the potential to change the physiological parameter of interest or together with use of an external energy source to change the parameter of interest.

Thus one example is to administer thermosensitive diagnostic agents in connection with an external heating and to follow the heating effect in parts of the body.

Another example is to administer compounds that change pH in connection with a pH-sensitive particulate diagnostic agent to follow the pH-profile in the area of interest.

Still another example is to cause the subject under study to inhale oxygen, after administration of an oxygen sensitive diagnostic agent, to follow oxygen uptake in tissue.

Early diagnosis is very important to obtain good therapeutic results. In most disease processes changes in physiological parameters take place before changes in morphology. All existing contrast agents diagnose morphology. The new types of contrast agent according to the invention are able to detect diseases at a very early stage in the disease process and thereby improve the therapeutic outcome for the patient.

Where the particulate diagnostic agent or a component thereof carries an overall charge, it will conveniently be used in the form of a salt with a physiologically acceptable counterion, for example an ammonium, substituted ammonium, alkali metal or alkaline earth metal cation or an anion deriving from an inorganic or organic acid. In this regard, meglumine salts are particularly preferred.

The diagnostic agents of the present invention may be formulated in conventional pharmaceutical or veterinary parenteral administration forms, e.g. suspensions, dispersions, etc., for example in an aqueous vehicle such as water for injections.

Such compositions may further contain pharmaceutically acceptable diluents and excipients and formulation aids, for example stabilizers, antioxidants, osmolality adjusting agents, buffers, pH adjusting agents, etc.

Where the agent is formulated in a ready-to-use form for parenteral administration, the carrier medium is preferably isotonic or somewhat hypertonic.

Where the particulate agent comprises a chelate or salt of an otherwise toxic metal species, e.g. a heavy metal ion, it may be desirable to include within the formulation a slight excess of a chelating agent, e.g.

- 44 -

as discussed by Schering in DE-A-3640708, or more preferably a slight excess of the calcium salt of such a chelating agent.

The dosage of the diagnostic agents of the invention will depend upon the imaging modality, the contrast generating species and the means by which contrast enhancement occurs (e.g. with switching on or off of contrast, with dispersion of contrast out of the vascular space, etc).

In general however dosages will be between 1/10 and 10 times the dosage conventionally used for the selected contrast generating species or analogous species in the same imaging modality. Even lower doses may also be used.

While the present invention is particularly suitable for methods involving parenteral administration of the particulate material, e.g. into the vasculature or directly into an organ or muscle tissue, intravenous administration being especially preferred, it is also applicable where administration is not via a parenteral route, e.g. where administration is transdermal, nasal, sub-lingual or is into an externally voiding body cavity, e.g. the gi tract, the bladder, the uterus or the vagina. The present invention is deemed to extend to cover such administration.

The disclosures of all the documents mentioned herein are incorporated by reference.

The present invention will now be illustrated further by reference to the following non-limiting Examples.

Example 1

Preparation of Temperature Sensitive Paramagnetic Liposomes

Liposomes containing GdHPDO3A (ProHance, Bracco Spa, Milan, Italy) and GdDTPA-BMA (Omniscan, Nycomed Amersham Imaging AS, Oslo, Norway) were prepared by the

thin film hydration method. Two different saturated phospholipid blends were used; one consisting of hydrogenated phosphatidyl choline (HPC) (Lipoid GmbH, Ludwigshafen, German) and hydrogenated phosphatidylserine-sodium (HPS) (NOF Corporation, Amagasaki, Japan); the other composed of DPPC and DPPG-sodium (Sygena Ltd, Liestal, Switzerland). The phospholipid mixtures contained 5% or 10% (w/w) of the negatively charged HPS and DPPG components. The phospholipid mixtures were dissolved in a chloroform/methanol mixture and the organic solution was evaporated to dryness under reduced pressure. DPPC/DPPG liposomes were formed by hydrating the lipid film with a pre-heated (55°C) aqueous solution (pH 7.4) of 250 mM GdDTPA-BMA or 250 mM Gd HPD03A. The HPC/HPS liposomes were prepared analogously but with a lipid hydration temperature of 70°C. The DPPC/DPPG and HPC/HPS liposomes were allowed to swell for 2 hours at 55 and 70°C respectively. The total lipid concentration was 50 mg/ml. The liposomes were subjected to 3 freeze-thaw cycles in liquid nitrogen. Differently sized liposomes were produced by sequential extrusion (Lipex Extruder[®], Lipex Biomembranes Inc., Vancouver, Canada) through polycarbonate filters of various pore diameters. The extrusion temperature was 55 and 70°C for the DPPC/DPPG and HPC/HPS liposomes respectively. Untrapped metal chelate was removed by gel filtration or dialysis against isoosmotic and isoprotic glucose solution.

Physiochemical Properties

The mean hydrodynamic diameter of the liposomes varied from 103 nm to 276 nm, as measured by photon correlation spectroscopy (ZetaSizer IV, Malvern Instruments Ltd., Malvern, England); the zeta potential was negative in the order of -25 mV, as determined by laser Doppler velocimetry at 25°C (ZetaSizer IV, Malvern Instruments Ltd., Malvern, England). The mean

gel-to-liquid crystalline phase transition temperature (T_c) of the HPC/HPS and DPPC/DPPG preparations was 50 and 42°C, respectively, as determined by differential scanning calorimetry (DSC4, Perkin Elmer Inc., Norwalk, CT).

Temperature Response of *In Vitro* MR Contrast Efficacy

Figure 1 of the accompanying drawings and Table 5 below show the temperature sensitivity of *in vitro* T_1 relaxivity (r_1) for liposome encapsulated GdDTPA-BMA and GdHPD03A, respectively (0.47T). Figure 2 of the accompanying drawings shows the temperature response of the *in vitro* MR signal intensity for liposome encapsulated GdDTPA-BMA.

Figure 3 of the accompanying drawings shows a series of T_1 -w GRE images prior to and after heating of a gel phantom containing inserts of liposome encapsulated GdDTPA-BMA.

Table 5

Temperature (°C)	r_1 ($s^{-1} mM^{-1}$)		
	DPPC/DPPG 103 nm	HPC/HPS 130 nm	Control
20	0.16	0.06	4.53
25	0.23	0.08	4.27
30	0.31	0.12	3.94
37	0.69	0.21	3.75
45	3.30	0.53	3.07
55	3.10	3.00	2.82
60	-	2.96	2.54

non-liposomal GdHPD03A

Example 2

GdDTPA-BMA encapsulated within DSPC/DPPC/DPPG liposomes

DSPC/DPPC/DPPG (weight ratio; 28.5/66.5/5)

liposomes were prepared by the thin film hydration method. The phospholipids (500 mg) were dissolved in a chloroform/methanol mixture and the organic solution was evaporated to dryness under reduced pressure. Liposomes were formed by hydrating the lipid film with a pre-heated (57°C) aqueous solution (pH \approx 7) of 250 mM GdDTPA-BMA (10 ml). The liposomes were subjected to 3 freeze-thaw cycles and allowed to swell for one and a half hours at 65°C. The liposome dispersion was extruded at 65°C through polycarbonate filters of various pore diameters. The liposome size (z-average) after extrusion was 167 nm. Untrapped GdDTPA-BMA was removed by dialysis against isoosmotic and isotonic glucose solution.

Table 6 shows the temperature sensitivity of the *in vitro* r_1 (0.235T) in glucose 5% solution for liposome encapsulated GdDTPA-BMA.

Table 6

Temperature (°C)	r_1 in glucose 5% (s ⁻¹ mM ⁻¹)
30	0.098
35	0.13
38	0.22
40	0.27
41	0.31
43	1.10
45	2.92

Example 3

GdDTPA-BMA encapsulated within DPPC/DPPG/DPPE-PEG-2000 liposomes

DPPC/DPPG/DPPE-PEG-2000 (weight ratio; 90/5/5) liposomes were prepared by the thin film hydration method. The phospholipids (500 mg) were dissolved in a chloroform/methanol mixture and the organic solution was

evaporated to dryness under reduced pressure. Liposomes were formed by hydrating the lipid film with a pre-heated (57°C) aqueous solution (pH \approx 7) of 250 mM GdDTPA-BMA (10 ml). The liposomes were subjected to 3 freeze-thaw cycles and allowed to swell for one and a half hours at 65°C. The liposome dispersion was extruded at 65°C through polycarbonate filters of various pore diameters. The liposome size (z-average) after extrusion was 132 nm. Untrapped GdDTPA-BMA was removed by dialysis against isoosmotic and isoprotic glucose solution.

Table 7 shows the temperature sensitivity of the *in vitro* r_1 (0.235T) in glucose 5% solution for liposome encapsulated GdDTPA-BMA.

Table 7

Temperature (°C)	r_1 in glucose 5% (s ⁻¹ mM ⁻¹)
35	0.32
37	0.46
38	0.56
39.2	2.53
40	4.16
42	5.65

Example 4

GdDTPA-BMA encapsulated within DSPC/DPPC/DPPG liposomes

DSPC/DPPC/DPPG (weight ratio; 43/52/5) liposomes were prepared by the thin film hydration method. The phospholipids (500 mg) were dissolved in a chloroform/methanol mixture and the organic solution was evaporated to dryness under reduced pressure. Liposomes were formed by hydrating the lipid film with a pre-heated (63°C) aqueous solution (pH \approx 7) of 250 mM GdDTPA-BMA (10 ml). The liposomes were subjected to 3 freeze-thaw cycles and allowed to swell for one and a half

hours at 64°C. The liposome dispersion was extruded at 65°C through polycarbonate filters of various pore diameters. The liposome size (z-average) was 145 nm. Untrapped metal chelate was removed by dialysis against isoosmotic and isotonic glucose solution.

Table 8 shows the temperature sensitivity of the *in vitro* r (0.235T) in both glucose 5% solution and human serum for liposome encapsulated GdDTPA-BMA.

Table 8

Temperature (°C)	r in glucose 5% (s ⁻¹ mM ⁻¹)	r in serum (s ⁻¹ mM ⁻¹)
35	0.12	0.14
40	0.22	0.25
42	0.29	0.44
44	0.88	1.91
46	4.47	4.51
48	4.40	4.51
50	4.40	4.35

Example 5

GdDTPA-BMA encapsulated within DPPC/DPPG liposomes

DPPC/DPPG (weight ratio; 95/5) liposomes were prepared by the thin film hydration method. The phospholipids (500 mg) were dissolved in a chloroform/methanol mixture and the organic solution was evaporated to dryness under reduced pressure. Liposomes were formed by hydrating the lipid film with a pre-heated (52°C) aqueous solution (pH \approx 7) of 250 mM GdDTPA-BMA (10 ml). The liposomes were subjected to 3 freeze-thaw cycles and allowed to swell for one and a half hours at 55°C. The liposome dispersion was extruded at 62°C through polycarbonate filters of various pore diameters. The liposome size (z-average) after extrusion was 148 nm. Untrapped metal chelate was removed by dialysis against isoosmotic and isotonic glucose

solution.

Table 9 shows the temperature sensitivity of the *in vitro* r (0.235 T) in both glucose 5% solution and human serum for liposome encapsulated GdDTPA-BMA.

Table 9

Temperature (°C)	r in glucose 5% (s ⁻¹ mM ⁻¹)	r in serum (s ⁻¹ mM ⁻¹)
35	0.331	0.389
38	0.753	0.810
39	1.47	1.20
40	3.75	3.31
41	4.88	5.05
42	4.80	4.99
44	4.80	4.78
48	4.77	4.88

Example 6

"Double transition" with a mixture of DSFC/DPPC/DPPG and DPPC/DPPG liposomes, containing both GdDTPA-BMA

1.5 ml liposomes from Example 4 were mixed with 1.5 ml DPPC/DPPG liposomes prepared as Example 5. The mixture was diluted to 40 ml with glucose 5% solution.

Table 10 shows the temperature sensitivity of the *in vitro* R_1 (0.235 T) in glucose 5% solution for the liposome mixture.

Table 10

Temperature (°C)	R in glucose 5% (s ⁻¹)
35	2.46
38	2.61
39	2.83
40	3.87
41	7.11
42	7.17
44	10.9
46	14.0
48	14.0

Example 7Perfluorobutane bubbles stabilised by 5mg/ml
DSPC/DPPC/DPPG

DSPC/DPPC/DPPG (weight ratio; 28.5/66.5/5) perfluorobutane gas bubbles were prepared by the thin film hydration method. The phospholipids (500 mg) were dissolved in a chloroform/methanol mixture and the organic solution was evaporated to dryness under reduced pressure. The lipid film was hydrated for 1 hour at 60°C after addition of 100 ml 1.5 % propylene glycol in water. The final dispersion contained 5 mg lipids/ml.

Five 2 ml vials were filled with 1 ml of the dispersion. The headspace was flushed with perfluorobutane gas. The vials were shaken on a CapMixer for 45 seconds and left on a roller table over night. The content of the five vials were collected and centrifuged for 5 minutes at 2000 rpm. The infranatant was removed and replaced by the same volume water. The microbubbles were reconstituted by gentle handshaking after flushing the headspace with perfluorobutane gas. The washing procedure was repeated three times.

The sample of perfluorobutane bubbles was characterized using Coulter Multisizer II fitted with an aperture of 50 μ m and Nycomed in-house equipment for measuring acoustic attenuation. The dispersion showed a size distribution of volume median diameter of about 3 μ m. The bubbles showed a nice attenuation spectrum in the range 3.5-8.0 MHz and were tested for pressure stability at an over-pressure of 150 mm Hg in the temperature range 22-47°C using the acoustic technique. The acoustic measurements showed that the gas bubbles disrupted at an over-pressure of 150 mm Hg at 47°C, whereas they remained stable at 40°C. This indicates that the gas microbubbles can be used in ultrasound imaging for in vivo mapping of physiological pressure.

Example 8

Imaging studies in rats with GdDTPA-BMA encapsulated within DPPC/DPPG liposomes

a) Intramuscular injection in the left thigh

Liposomes were injected intramuscularly at a dosage of 0.02 mmol/kg. The left thigh muscle was heated with focused ultrasound whereas the right thigh muscle served as a control.

Figures 4-5 show axial T-w SE images of the thigh before and after liposome injection, respectively. Figures 6-8 are T-w SE images after 2, 5, and 9 minutes of heating, respectively.

At that timepoint, heating was terminated, the rat was removed from the MRI scanner and the temperature of the muscle was measured to be 47°C. Figure 9 represents the final image 15 minutes after termination of heating. For comparative purposes, the syringe containing the liposomal dispersion (identical to that injected) was included.

The results indicate that the signal intensity of the left thigh muscle increases substantially after heating, as compared to the right thigh muscle and syringe.

b) Intravenous injection

Liposomes were injected intravenously into a rat (upper position) at a dosage of 0.10 mmol/kg. The rat in the lower position served as control (e.g. no injection nor heating).

Figure 10 is the axial T-w SE image of the liver 7 minutes after liposome injection. At 15 minutes post injection, the liver was heated by focused ultrasound (Figure 11). Figures 12-13 are T-w SE images 16 and 21 minutes after initiation of heating, respectively.

After termination of heating, the measured temperature in the liver was 51°C.

The results indicate that the liver signal intensity increases substantially after heating as compared to the control liver.

Example 9

Preparation of pH-sensitive paramagnetic liposomes

Liposomes composed of DPPE/PA (4:1 mol/mol) containing GdDTPA-BMA were prepared by the thin film hydration method. The total lipid concentration was 25 mg/ml. Briefly, a chloroform/methanol (10:1) solution of the lipids was rotary evaporated to dryness and the resulting film was further dried under vacuum over night. The lipids were hydrated with 250 mM GdDTPA-BMA in 0.05 M Tris-HCl buffer (pH = 8.4) at 75 °C. The liposomes were subjected to 3 freeze-thaw cycles in MeOH/CO₂(s). The liposomes were sized down by sequential extrusion (Lipex Extruder[®], Lipex Biomembranes Inc., Vancouver, Canada) through polycarbonate filters with various pore diameters. Untrapped metal chelate was removed by dialysis against isoosmotic glucose solution (pH=8.4).

Physicochemical properties

The mean hydrodynamic diameter of the liposomes was measured to 165 nm by photon correlation spectroscopy (ZetaSizer IV, Malvern Instruments Ltd., Malvern, England). The *in vitro* T₁-relaxation times of the paramagnetic liposomes were measured (0.235 T, Minispec PC-110b, Bruker GmbH, Rheinstetten, Germany) in different isoosmotic buffer solutions (0.05 M citrate-phosphate buffer and 0.05 M Tris-HCl buffer). The investigated pH range was 4-8.5. The buffered liposome dispersions were incubated at 37°C for 15 minutes. Table 11 shows the pH sensitivity of *in vitro* r₁-relaxivity for liposome encapsulated GdDTPA-BMA.

Table 11

pH dependency of the r (37 °C, 0.235 T) for liposomal GdDTPA-BMA

pH	r (s ⁻¹ mM ⁻¹)
3.91	1.32
4.30	1.26
4.70	1.31
5.15	1.17
5.59	1.10
5.95	1.03
6.40	1.00
6.71	0.50
7.33	0.32
7.69	0.29
8.02	0.28
8.34	0.29
8.54	0.31

Example 10

DyDTPA-BMA encapsulated within DPPC/DPPG liposomes

DPPC/DPPG (weight ratio; 95/5) liposomes were prepared by the thin film hydration method. The phospholipids (500mg) were dissolved in a chloroform/methanol mixture and the organic solution was evaporated to dryness under reduced pressure. Liposomes were formed by hydrating the lipid film at 50°C with an aqueous solution (pH \approx 7) of 250 mM DyDTPA-BMA (sprodiamide, Nycomed Imaging AS, Oslo, Norway) (10ml). The liposomes were subjected to 3 freeze-thaw cycles and allowed to swell for one hour at 59°C. The liposome dispersion was extruded at 65°C through polycarbonate filters of various pore diameters. The liposome size (z-average) after extrusion was 153 nm. Untrapped DyDTPA-BMA was removed by dialysis against isoosmotic and isotonic glucose solution. The temperature sensitivity of the MR contrast effect may be investigated.

Example 11GdDTPA-dextran encapsulated within DPPC/DPPG liposomes

DPPC/DPPG (weight ratio; 95/5) liposomes were prepared by the thin film hydration method. The phospholipids (500 mg) were dissolved in a chloroform/methanol mixture and the organic solution was evaporated to dryness under reduced pressure. The liposomes were formed by hydrating the lipid film at 48°C with an aqueous solution of 50 mM GdDTPA-dextran (MW 156 kD), whose synthesis is described in: P Rongved et al., Carbohydr. Res., 287 (1996) 77-89. The liposome dispersion was sonicated at 46°C using a sonicator tip. The liposome size (z-average) after sonication was 70 nm. Untrapped GdDTPA-dextran is removed by gel filtration or dialysis against isoosmotic and isoprotic glucose solution. The temperature sensitivity of the MR contrast effect may be investigated.

Example 12GdDTPA-BMA encapsulated within dibehenoyl-PC liposomes

Dibehenoyl-PC (22:0) (Table 1) liposomes may be prepared by the thin film hydration method. The phospholipids (500 mg) are dissolved in a chloroform/methanol mixture and the organic solution is evaporated to dryness under reduced pressure. Liposomes are formed by hydrating the lipid film at 80°C with an aqueous solution (pH \approx 7) of 250 mM GdDTPA-BMA (10 ml). The liposomes are subjected to 3 freeze-thaw cycles and allowed to swell for one and half-hours at 80°C. The liposome dispersion is extruded at 80°C through polycarbonate filters of various pore diameters. Untrapped GdDTPA-BMA is removed by gel filtration or dialysis against isoosmotic and isoprotic glucose solution. The temperature sensitivity of the MR contrast effect may be investigated.

Example 13Superparamagnetic iron oxides (SPIOs) encapsulated within HPC/HPS liposomes

HPC/HPS (weight ratio; 90/10) liposomes were prepared by a modified thin film hydration method. Liposomes were formed by adding a homogeneous mixture of phospholipids (700 mg) to 10 ml of a pre-heated (55 C) aqueous dispersion of PEGylated SPIOs (6.10 mg iron/ml). The liposomes were allowed to swell for 30 minutes at 65 C. The liposome dispersion was extruded at 66 C through polycarbonate filters of various pore diameters. Untrapped SPIOs are removed by gel filtration or dialysis. The temperature sensitivity of the MR contrast effect may be investigated.

Example 14Ultrasmall superparamagnetic iron oxides (USPIOs) encapsulated within HPC/HPS liposomes

HPC/HPS (weight ratio; 90/10) liposomes were prepared by a modified thin film hydration method. Liposomes were formed by adding a homogeneous mixture of phospholipids (700 mg) to 10 ml of a pre-heated (70°C) aqueous dispersion of USPIOs (3.63 mg iron/ml). The liposomes were allowed to swell for 90 minutes at 70°C. The liposome dispersion was extruded at 70°C through polycarbonate filters of various pore diameters. Untrapped USPIOs are removed by gel filtration or dialysis. The temperature sensitivity of the MR contrast effect may be investigated.

Example 15Superparamagnetic iron oxides (SPIOs) or ultrasmall-SPIOs encapsulated within pH-sensitive liposomes

SPIOs or USPIOs encapsulated within pH-sensitive

- 58 -

liposomes may be prepared in a manner analogous to that used in Example 9. Untrapped superparamagnetic material is removed by gel filtration or dialysis. The pH-sensitivity of the MR contrast effect may be investigated.

Example 16

GdDTPA-BMA encapsulated within DSPC/DMPG/cholesterol liposomes

DSPC/DMPG/cholesterol liposomes (molar ratio; 49:5:20) were prepared by a modified thin film hydration method. Liposomes were formed by adding a freeze-dried mixture of phospholipids (60 g) to a pre-heated (59°C) aqueous solution (pH \approx 6.3) of 250 mM GdDTPA-BMA/300 mM sucrose/10mM phosphate (300 ml). The liposomes were allowed to swell for 30 minutes at 59°C. The liposome dispersion was homogenized and extruded at high pressure through polycarbonate filters with a pore size of 400 nm. Untrapped GdDTPA-BMA was removed by ultrafiltration with a 300 mM sucrose/10 mM phosphate solution. The liposome size (z-average) after ultrafiltration was 110 nm. Liposomes were also lyophilized (2ml per vial) and reconstituted by addition of 2 ml of deionized water. The liposome size (z-average) after reconstitution was 119 nm.

Table 12 summarizes the temperature sensitivity of the in vitro R_1 (0.235 T) for liposome encapsulated GdDTPA-BMA in a 300 mM sucrose/10 mM phosphate solution. The influence of lyophilization on the R_1 -temperature sensitivity of the reconstituted liposomes is also shown in Table 12.

Table 12

	T Relaxation Time (ms)		
	35°C	40°C	55°C
Before lyophilisation	930	750	350
After lyophilisation	670	590	340

Example 17Commercially available Gd compounds/Gd compounds in development phase encapsulated within temperature- or pH-sensitive liposomes

Liposomes containing the following contrast agents: GdBOPTA (Bracco spa, Italy), GdDTPA (Shering AG, Berlin), GdDOTA (Guerbet SA, Aulnay-sous-Bois), Gadomer (Shering AG, Berlin) MS-325 and protein bound MS-325 (Epix Medical Inc, USA) may be prepared in a manner analogous to that used in Examples 1-5, 9 and 12. Untrapped Gd compound is removed by gel filtration or dialysis. The temperature sensitivity of the MR contrast effect may be investigated.

Example 18Temperature sensitivity of *in vitro* r, in blood for liposome encapsulated GdDTPA-BMA

DPFC/DPPG/DPPE-PEG-2000 and DPFC/DPPG liposomes containing GdDTPA-BMA were prepared in a manner analogous to that used in Examples 3 and 5, respectively. Table 13 summarizes the temperature evolution of the *in vitro* r (0.235 T) in rat blood for both liposome formulations.

Table 13

Temperature (°C)	r_1 ($s^{-1}mM^{-1}$) DPPC/DPPG 120 nm	r_1 ($s^{-1}mM^{-1}$) DPPC/DPPG/DPPE- PEG 121 nm
35	0.260	0.244
37	0.479	0.391
39	0.659	0.588
40	1.18	0.823
41	2.45	1.26
42	4.06	2.87
43	5.18	3.65
44	4.57	4.06

Example 19

Pilot biodistribution and relaxometric studies of
liposomal GdDTPA-BMA in male rats

a) Intramuscular injection

DPPC/DPPC liposomes containing GdDTPA-BMA (prepared in Example 18) were injected intramuscularly (im) into Sprague Dawley rats at a dosage of 20 :mol/kg.

Table 14 shows the T_1 relaxation times (37°C, 0.235 T) of excised tissues and blood one and three hours after im injection of DPPC/DPPG liposomes (n= 2x3). Table 15 shows the temperature response of the T_1 in muscle. All results are given as mean values.

Table 14

Time post Injection (min)	T ₁ relaxation time (ms)		
0	500	833	248
60	451	950	249
180	440	833	243

Table 15

Time post Injection (min)	Muscle T ₁ (ms)	
0	500	340
60	451	341
180	440	383

Despite large interindividual variations, the results show the temperature dependence of the muscle T₁ after im administration of liposome encapsulated GdDTPA-BMA.

b) Intravenous injection

DPPC/DPPG/DPPE-PEG-2000 and DPPC/DPPG liposomes containing GdDTPA-BMA (prepared in *Example 18*) were injected intravenously (iv) into Sprague Dawley rats at a dosage of 100 :mol/kg.

Tables 16 and 17 show the T relaxation times (37°C,

- 62 -

0.235 T) of excised tissues and blood, 5 minutes (n=3), one (n=2) and three (n=3) hours after iv injection of DPPC/DPPG and DPPC/DPPG/DPPE-PEG-2000 liposomes, respectively. Also shown, is the Gd uptake in tissue, expressed as the percentage tissue uptake of the administered Gd dosage. Tables 18 and 19 summarize the temperature response of the blood T after iv injection of DPPC/DPPG and DPPC/DPPG/DPPE-PEG-2000 liposomes, respectively. A more detailed investigation of the temperature response was performed in blood withdrawn one hour after iv administration (n=3) of DPPC/DPPG liposomes, as shown in Table 20. All results are given as mean values.

Table 16

Time post injection (min)	T relaxation time (ms)			
	<u>Tissue uptake (% of adm. Gd dosage)</u>			
	Liver	Blood	Spleen	Lungs
0	248	833	510	613
	-	-	-	-
5	212	533	357	547
	11.6	<u>60.0</u>	<u>1.9</u>	<u>0.88</u>
60	223	589	315	520
	<u>13.7</u>	<u>25.9</u>	<u>4.3</u>	<u>0.48</u>
180	227	823	353	557
	<u>7.9</u>	<u>1.3</u>	<u>2.0</u>	<u>0.06</u>

Table 17

Time post injection (min)	T relaxation time (ms)			
	<u>Tissue uptake (% of adm. Gd dosage)</u>			
	Liver	Blood	Spleen	Lungs
0	248	833	510	613
	-	-	-	-
5	224	540	417	580
	<u>5.7</u>	<u>64.3</u>	<u>1.4</u>	<u>0.85</u>
180	202	500	298	563
	<u>8.8</u>	<u>27.7</u>	<u>6.5</u>	<u>0.56</u>

Table 18

Time post Injection (min)	Blood T ₁ (ms)	
	37°C	43°C
0	833	880
5	533	151
60	589	273
180	823	797

Table 19

Time post Injection (min)	Blood T (ms)	
	37°C	43°C
0	833	880
60	540	135
180	500	244

Table 20

Blood T (ms)			
37°C	40°C	41°C	43°C
520	247	232	210

The results show the potential of both non-PEGylated and, especially, PEGylated liposomal GdDTPA-BMA as blood pool agents. The T₁-temperature sensitivity of the liposomes was also demonstrated in blood.

Example 20

In Vitro Imaging Studies with GdDTPA-BMA encapsulated within DSPC/DPPC/DPPG liposomes

DSPC/DPPC/DPPG liposomes containing GdDTPA-BMA were prepared analogously to Example 2. The liposome size (z-average) was 129 nm. MR imaging was performed at 2.0 T (Bruker Medspec) on a concentric spherical phantom in which the inner chamber contained liposomal GdDTPA-BMA diluted with an isotonic medium composed of glucose and 6.25% polyvinylpyrrolidone (conc. \approx 0.8 mM Gd), whilst the outer compartment was filled with saline. Microwave heating was performed at 434 MHz with a linear radio

frequency antenna placed in the outer chamber. The microwave irradiation was applied simultaneously with the image acquisition. Blocks consisting of 10 diffusion-weighted spin-echo single shot EPI (DW-SE-EPI) images (b-factor from 3 to 864 s/mm), a set of SE-EPI images with inversion-recovery preparation (IR-SE-EPI) and gradient echo T₂-weighted (T₂-W-GE) images were repeated until the temperature of the liposome sample reached 48°C (in appr. 110 min). T₂-W-GE images were acquired with TE/TR/flip: 5ms/30ms/50°. T₂-maps were calculated from the set of 13 IR-SE-EPI images, measured with inversion times varying from 14.4 ms to 16 s. Plots of 1/T₂ (R₂) versus temperature were generated from a fixed region-of-interest within the phantom. The sample temperature was measured by a thermocouple immediately after acquisition of each block. The temperature distribution within the imaged slice was evaluated from ADC-maps.

The temperature evolution of the measured R₂ for liposomal GdDTPA-BMA is summarized in Figure 14. A linear correlation was obtained between R₂ and temperature in the "transition region" 40.4-43.7°C (regression coefficient of 0.995). Figure 15 shows selected T₂-W GE images of the phantom (a) before heating, (b) during heating; signal intensity distribution observed within liposome sample, and (c) after heating; homogeneous signal intensity distribution. Figure 16 shows the corresponding T₂-maps at the same time points as for Figure 15. By use of the linear correlation between R₂ and temperature, a corresponding temperature map could be derived from the T₂-map at timepoint (b), as seen on Figure 17. The temperature map demonstrates the thermosensitivity of liposomal GdDTPA-BMA. (NB The temperature scale is only valid for the inner chamber containing liposomes).

Example 21

In vitro MR imaging studies with GdDTPA-BMA encapsulated within DPPC/DPPG liposomes - Determination of Gd concentration.

A static *in vitro* phantom, composed of twelve glass vials (10 mm dia.) placed in a rectangular plastic container, was used for this study. The plastic container was filled with a viscous isotonic medium composed of glucose/25% (w/w) polyvinyl-pyrrolidone (PVP) and doped with GdDTPA-BMA to give a T_1 of about 430 ms at 1.5 T. Three of the vials contained a marker solution with a known T value (about 630 ms). The remaining nine vials were filled with DPPC/DPPG-based GdDTPA-BMA liposomes (prepared in Example 18) dispersed in varying amounts of isotonic 10% PVP/glucose solution. The Gd concentration [Gd] in the liposome samples ranged from 0 to 5.2 mM Gd as determined by inductively coupled plasma atomic emission spectrophotometry. The phantom was imaged at room temperature in a quadrature knee coil at 1.5 T on a Philips NT system. The following imaging sequences were used:

1. T_1 -FFE (spoiled gradient echo) TR/TE/flip: 15ms/2ms/30°.
2. TMIX (quantitative T_1/T_2 sequence)
3. Dual TE FFE (T_2^* mapping) (TE1/TE2: 4ms/50 ms)

All three sequences were repeated after heating of the phantom, the latter achieved by placing the phantom in a warm (>60°C) water bath. The temporal effect of heating was not investigated, only the end-effect (i.e. $T \gg T_1$).

A linear correlation was obtained between [Gd] and matrix corrected R_1 (ΔR_1) prior to heating (measured from TMIX sequence), with a regression coefficient of 0.996. The calculated liposomal r was $0.11 \text{ mM}^{-1}\text{s}^{-1}$.

- 67 -

Analogously, a liposomal r of 0.55 mM s was determined from the R vs $[Gd]$ curve (regression coefficient = 0.997); the r/r ratio being equal to 5. After liposome heating, the r and r were 3.23 and 3.75, respectively, giving an r/r ratio of 1.16.

Figure 18 shows, prior to heating, a linear correlation between $[Gd]$ and the ratio of the signal intensities of liposome sample and PVP gel (SI_{lip}/SI_{PVP}) using the T-FFE sequence. Figure 19 shows the plot of the (R_1/R_2) ratio vs the $[Gd]/[Gd]_{ref}$ ratio; here $[Gd]_{ref} = 5.2$ mM. The results demonstrated that prior to heating, the (R_1/R_2) ratio accurately reflected the $[Gd]$ ratio. Similar results were also obtained when the (R_1/R_2) ratio was employed. The findings suggest that the T₁- (and T₂-) effect of liposome encapsulated GdDTPA-BMA prior to heating is significant enough to enable a relative assessment of liposomal Gd concentration.

Example 22

In vitro MR imaging studies with GdDTPA-BMA encapsulated within DPPE/PA liposomes

DPPE/PA liposomes containing GdDTPA-BMA were prepared analogously to Example 9. The mean hydrodynamic diameter of the liposomes was measured to 158 nm by photon correlation spectroscopy (Malvern PS/MW 4700, Malvern Instruments Ltd., Malvern, England). An in vitro phantom, composed of thirteen glass vials (11 mm diameter) placed in a circular glass reactor, was used for this study. The glass reactor was filled with an agar gel (2 % w/v) doped with GdDTPA-BMA to give a T_1 of about 900 ms at 1.5 T. The glass vials were filled with isoosmotic buffer solutions with pHs ranging from 4.8 to 8.2. The phantom was constantly held at a temperature of 37°C by circulating heated water through the shell of the reactor with a circulating water pump. Liposomes were added successively to each vial with a

time interval of 1 minute. The imaging was started 25 minutes after addition of liposomes to the first vial. The phantom was imaged at 1.5 T on a Philips NT system. The following imaging parameters were used: sequence: MIX-TSE; TR (ms): 800.0; TE (ms): 12.5; TI (ms): 500.0; flip (deg): 90; slice thickness (mm): 7.0; FoV (freq*phase, mm): 230.0*230.0. The scan cycles were repeated every minute for 20 minutes. Figure 20 shows the phantom 25 minutes after addition of liposomes to the first vial. The signal intensity increases with decreasing pH.

Example 23

Particles from polymer made from ethylidene bis(16-hydroxyhexadecanoate) and adipoyl chloride

Air filled particles of the polymer ethylidene bis(16-hydroxyhexadecanoate) and adipoyl chloride were made as described in Example 3f of WO 96/07434.

Change of gas

The dry powder was exposed to 20 mmHg vacuum for ca. 15 minutes, followed by inlet of perfluorobutane gas. The powder of polymer particles containing perfluorobutane gas were then redispersed to 10 mg/ml dry material in MilliQ water by shaking on a laboratory shaker for 12-16 hours. Examination by light microscopy indicated formation of a particle dispersion with irregular shaped particles. The particles floated readily, as expected for gas-containing particles.

Heating of the polymer particles

10 ml of the polymer dispersion was heated on a water bath to 65°C in one minute while using magnet stirring. At this temperature the polymer melted. Microscopic evaluation indicated that the irregular particles changed to spherical, smooth particles,

indicating that the polymer capsule melted.

Characterization

The acoustic effect of the suspension prepared above was obtained by measuring the ultrasonic transmission through a dispersion in an aqueous carrier liquid, using a 3.5 MHz broadband transducer in a pulse-reflection technique. The aqueous carrier liquid was used as reference. Table 21 contains the observed acoustic attenuation compared to non-heated particle dispersion at the same concentrations, indicating that heat treatment removes most of the acoustic attenuation.

The gas content was measured by density measurements before and after destruction of the microcapsules using high-energetic ultrasound. The results show that the gas content is conserved. The results show that by melting the polymer capsules, the gas filled microcapsules become almost invisible to ultrasound, probably due to the stiff polymer shell now surrounding the gas phase.

Table 21. Characteristics of perfluorobutane containing polymer particles treated by heating the dispersions to 65°C.

Description	Gas content [% v/v (rel. sample vol.)]	Acoustic attenuation [dB/cm]
Reference	1.398	6.42
65°C, 1 min	1.227	0.63

Example 24

The particles from example 23 above are studied by acoustic characterisation in vitro by measuring the ultrasonic transmission through a dispersion in an aqueous carrier liquid, using a 3.5 MHz broadband

- 70 -

transducer in a pulse-reflection technique. The aqueous carrier liquid is used as reference. The acoustic characterisation is started at room temperature where the acoustic attenuation is low as shown in Table 21 above. The temperature is raised and acoustic characterisations are done at different temperature intervals. When the temperature passes 48.6°C, the melting point of the polymer (see example 23 above), the acoustic attenuation increases sharply, indicating a temperature sensitive contrast agent. A similar experiment could be done using a polymer with melting point around 37-40°C, and hence closer to body temperature.

Example 25

A spatula edge of micronised kaolin is added to 2 ml perfluorodimethylcyclobutane (b.p. 45°C) and dispersed using 0.2 ml Fluorad™ FC-171 surfactant. A milky white dispersion is obtained by vigorously stirring.

1 ml of a dispersion of 1,2-distearoyl-phosphatidyl glycerol and (0,5 mg/ml) and distearoylphosphatidyl-choline (4,5 mg/ml) in purified water is placed in a 2 ml vial to which is added 100 µl of the kaolin in perfluorodimethylcyclobutane dispersion described above. The vial is closed and then shaken for 75 seconds using an Espe CapMix® to yield a kaolin in perfluorodimethylcyclobutane in water emulsion.

An acoustic apparatus is mounted in which the acoustic effect of the suspension prepared above can be obtained by measuring the ultrasonic transmission through a dispersion in an aqueous carrier liquid, using a 3.5 MHz broadband transducer in a pulse-reflection technique. The aqueous carrier liquid can be used as reference. The sample is injected in a thermostatted cell, where an overpressure or an underpressure can be

- 71 -

applied by a pump.

The emulsion is transformed to the acoustic cell and diluted with 50 ml water, keeping the temperature constant at 37°C. The first measurement is done at atmospheric pressure showing weak acoustic attenuation due to the presence of liquid emulsion droplets and no gas bubbles in the cell. The pressure is then gradually reduced at intervals, and the acoustic measurement is done at each interval. When the pressure reach 580 mmHg (i.e. 180 mmHg below atmospheric pressure), the acoustic attenuation increases sharply and significantly, demonstrating that the microdroplets now boil and turn to acoustic effective microbubbles.

This experiment demonstrate how an emulsion with a disperse phase of boiling point slightly above body temperature can be used to map underpressure in vivo, for instance the underpressure which will occur below an embolisation of a blood vessel.

Example 26

GdDTPA-BMA encapsulated within Mg⁺⁺ sensitive liposomes

Beef-heart cardiolipin - cesium salt, dipalmitoylphosphatidylcholine (DPPC) and dipalmitoylphosphatidylglycerol - potassium salt (DPPG) are added to a round bottom flask in 40/55/5 mol ratio (totally 500 mg) and dissolved using chloroform. The chloroform is removed by evaporation under reduced pressure using a rotavapor. Liposomes are formed by hydrating the lipid film with a pre-heated (52°C) aqueous solution of 250 mM GdDTPA-BMA (10 ml). The liposomes are subjected to 3 freeze-thaw cycles and allowed to swell for one and half-hour at 55°C. The liposome dispersion is extruded at 62°C through polycarbonate filters of various pore diameters. Untrapped GdDTPA-BMA is removed by dialysis against isosmotic and isotropic glucose solution.

The liposome dispersion is diluted ten times with water and transformed to an NMR tube. The r_1 relaxivity at 0.235 Tesla is measured using a Minispec NMR instrument at 37 °C. The r_1 relaxivity is low. The sample is then titrated by a 0,6 M MgCl₂ solution. When the Mg²⁺ to cardiolipin ratio increases, the lamellar to H₂O phase transition is induced as described in F. Reiss-Husson, *J. Mol. Biol.*, **25**, 363, (1967). The break-down of the liposomal structure leads to contact between the GdDTPA-BMA and water, inducing a significant increase in the r_1 relaxivity. This experiment will demonstrate a Mg²⁺ sensitive MRI contrast agent.

Example 27

GdDTPA-BMA encapsulated within Ca²⁺ sensitive liposomes

The experiment as described in Example 26 above is repeated, but the MgCl₂ solution is replaced by a CaCl₂ solution. Observations of a similar increase in relaxivity at a sufficiently high Ca²⁺ concentration demonstrate a Ca²⁺ sensitive MRI contrast agent.

Example 28

Perfluorobutane microbubbles stabilised by Phosphatidylserine as example of a Ca²⁺ sensitive ultrasound contrast agent

Preparation

Hydrogenated phosphatidylserine (5 mg/ml in a 1% w/w solution of propylene glycol in purified water) and perfluorobutane gas were homogenised by an Ystral® rotor-stator at 7800 rpm and ca. 40°C to yield a creamy-white microbubble dispersion. The dispersion was fractionated to substantially remove undersized microbubbles (<2 µm) and the volume of the dispersion was adjusted to the desired microbubble concentration by adding aqueous sucrose to give a sucrose concentration

- 73 -

of 92 mg/ml. 2 ml portions of the resulting dispersion were filled into 10 ml flat-bottomed vials specially designed for lyophilisation, and the contents were lyophilised to give a white porous cake. The lyophilisation chamber was then filled with perfluorobutane and the vials were sealed. Prior to use, water was added to a vial and the contents were gently hand-shaken for several seconds to give a perfluorobutane microbubble dispersion.

Microscopic investigation

One drop of the microbubble dispersion was placed on an object glass for microscopy investigation. The sample was covered with a cover glass and placed under a microscope. Droplets of a 50 mg/ml calcium chloride solution in water were added to the edge of the cover glass so that the solution penetrated into the microbubble dispersion. The behaviour of the microbubble dispersion as the calcium chloride solution front moved was recorded on videotape. Microbubble aggregates with larger dimensions than the initial microbubbles were observed to form, demonstrating that microbubbles with potential to have different ultrasound properties were generated.

Example 29

TcDTPA encapsulated within DPPC/DPPG liposomes

TcDTPA is prepared from sodium pertechnetate and a commercial kit containing SnCl₄ and DTPA. TcDTPA is encapsulated in liposomes similar to Example 10 above. The product is a temperature sensitive contrast agent for scintigraphic studies.

Example 30Gadolinium DTPA labelled starch microspheres

Gadolinium DTPA starch particles were prepared according to P. Rongved *et al.* in Carbohydrate Research 214 (1991) 325-330 substrate 9 to 12. The particles were suspended in 0.9 % NaCl solution before administration. The product can be used to diagnose diseases related to abnormal enzyme activity (α -amylase and esterase); for example.

Example 31Iodixanol-containing liposomes

A diagnostic composition comprising:

Iodixanol (total amount)	400 mg/ml
Iodine encapsulated	80 mg/ml
Sorbitol	20 mg/ml
Trometamol (TRIS)	0.097 mg/ml
EDTANa ₂ Ca	0.1 mg/ml
Hydrogenated phosphatidylcholine	51.2 mg/ml
Hydrogenated phosphatidylserine	5.1 mg/ml
Water for injection ad	1 ml (approx 0.9 ml)

was prepared by dissolving the phospholipid in chloroform/methanol/water (4:1:0.025, volume) and evaporating the solvent (rotary evaporation). An isotonic solution of iodixanol and sorbitol was made and heated to 60-70°C and this temperature was maintained during the remainder of the process. The phospholipid mixture was added with stirring, and the liposomes were formed. To control the size of the liposomes the preparation was homogenized (Rotor/Stator homogenizer). The liposomes were then extruded through 7 polycarbonate filters placed in series (pore diameter 1 μ m). The

- 75 -

product was diluted with an isotonic solution of iodixanol and sorbitol, and trometamol and EDTA were added. The product was filled into glass vials and autoclaved (121°C, 15 minutes). Tc = 49 °C.

The product can be used to monitor temperature during hyperthermia treatment with focused ultrasound.

Example 32

Gas containing microbubbles of DPPC/PPG/DPPE-PEG "doped" with vector with affinity for angiogenesis

Various products can be prepared using the technology described in WO 98/18500. The products can be used as tumor specific markers for hyperthermia treatment.

Claims

1. A method of imaging of an animate human or non-human animal body, which method comprises: administering parenterally to said body a particulate material comprising a matrix or membrane material and at least one contrast generating species, which matrix or membrane material is responsive to a pre-selected physiological parameter whereby to alter the contrast efficacy of said species in response to a change in the value of said parameter; generating image data of at least part of said body in which said species is present; and generating therefrom a signal indicative of the value or variation of said parameter in said part of said body.

2. A method as claimed in claim 1 wherein the physiological parameter is pH, temperature, pressure, carbon dioxide tension, enzyme activity, tissue electrical activity, tissue diffusion or ion concentration.

3. A method as claimed in claim 2 wherein the physiological parameter is pH, temperature or pressure.

4. A method as claimed in any one of claims 1 to 3 wherein the response of the matrix or membrane material to a change in the value of the pre-selected physiological parameter is a change in matrix or membrane permeability or chemical or physical breakdown of the matrix or membrane material.

5. A method as claimed in any one of claims 1 to 4 wherein the imaging technique is MRI, scintigraphy or ultrasound or X-ray imaging.

- 77 -

6. A method of MRI as claimed in claim 5 wherein the contrast generating species is a paramagnetic and/or superparamagnetic compound and/or an iron oxide or a gadolinium or dysprosium compound.
7. A method of ultrasound imaging as claimed in claim 5 wherein the contrast generating species is an encapsulated gas selected from air, a fluorohydrocarbon, sulphur hexafluoride and a perfluorocarbon.
8. A method of ultrasound imaging as claimed in claim 5 wherein the particulate material comprises a temperature, pressure or pH sensitive emulsion or suspension.
9. A method as claimed in claim 1 wherein said particulate material is in combination with a targeting ligand for a cell or receptor of interest.
10. A method as claimed in claim 1 wherein the membrane material forms a vesicle.
11. A method as claimed in claim 1 wherein the matrix or membrane material is selected from a phospholipid and a physiologically acceptable polymer.
12. A method as claimed in claim 10 or 11 wherein the membrane material forms a temperature or pH sensitive liposome.
13. A method as claimed in claim 12 wherein the liposome is stable at normal body temperature but exhibits increased water permeability or leakage at temperatures greater than normal body temperature.
14. A method as claimed in claim 1 wherein the contrast efficacy is altered by interaction between the contrast

generating species and the environment in the part of the animal body where the matrix or membrane material has responded to a change in the value of the physiological parameter.

15. A method as claimed in any one of the preceding claims wherein the physiological parameter is temperature and wherein the change in the value of said parameter is related to cancer, cardiovascular disease or inflammation or results from the treatment of hyperthermia in the animal body.

16. A method as claimed in any one of claims 1 to 14 wherein the physiological parameter is pH and wherein the change in the value of said parameter is caused by cancer, cardiovascular disease, osteoporosis, inflammations or autoimmune diseases.

17. A method as claimed in any one of claims 1 to 16 wherein in addition to the generation of a signal indicative of the value or variation of a pre-determined physiological parameter in a part of the animal body in which the contrast generating species is present, an anatomical image of the same part of the animal body is generated.

18. A method as claimed in claim 17 wherein no contrast agent is used to generate the anatomical image.

19. A method as claimed in claim 17 wherein a contrast agent is used in the generation of the anatomical image.

20. A method as claimed in claim 19 wherein the same contrast agent is used to generate a signal relating to the pre-selected physiological parameter and the anatomical image.

- 79 -

21. A contrast medium for imaging of a physiological parameter, said medium comprising a particulate material the particles whereof comprise a matrix or membrane material and at least one contrast generating species, said matrix or membrane material being responsive to said physiological parameter to cause the contrast efficacy of said contrast generating species to vary in response to said parameter.

22. The use of a contrast generating species for the manufacture of a particulate contrast medium for use in a method of diagnosis comprising generating a signal indicative of the value of said physiological parameter, the particles of said contrast medium comprising a matrix or membrane material and at least one contrast generating species, said matrix or membrane material being responsive to said physiological parameter to cause the contrast efficacy of said contrast generating species to vary in response to said parameter.

23. A method of imaging of an animate human or non-human animal body, which method comprises:

administering parenterally to said body at least one contrast generating species the contrast efficacy whereof is responsive to a change in value of a pre-selected physiological parameter;

generating image data of at least part of said body in which said species is present; and

generating therefrom a signal indicative of the value or variation of said parameter in said part of said body and also generating an anatomical image of the same part of the animal body.

ABSTRACT OF THE DISCLOSURE

The present invention relates to a method of imaging of an animate human or non-human animal body, which method comprises: administering parenterally to said body a particulate material comprising a matrix or membrane material and at least one contrast generating species, which matrix or membrane material is responsive to a pre-selected physiological parameter whereby to alter the contrast efficacy of said species in response to a change in the value of said parameters; generating image data of at least part of said body in which said species is present; and generating therefrom a signal indicative of the value or variation of said parameter in said part of said body. The invention also relates to contrast media for imaging a physiological parameter.

100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224
2225
2226
2227
2228
2229
2230
2231
2232
2233
2234
2235
2236
2237
2238
2239
2240
2241
2242
2243
2244
2245
2246
2247
2248
2249
2250
2251
2252
2253
2254
2255
2256
2257
2258
2259
2260
2261
2262
2263
2264
2265
2266
2267
2268
2269
2270
2271
2272
22





FIG. 1

TEMPERATURE RESPONSE OF IN VITRO r_1 FOR GdDTPA-BMA
ENCAPSULATED IN DPPC/DPPG LIPOSOMES (0.47T)

2 / 13

**FIG. 2**

TEMPERATURE RESPONSE OF MR SIGNAL INTENSITY FOR GdDTPA-BMA
ENCAPSULATED WITHIN DPPC/DPPG LIPOSOMES (2.0 T).




Figure 3. Gel phantom (a) containing inserts of DPPC/DPPG-based GdDTPA-BMA liposomes (labelled 3-10) and control glucose 5% solution (labelled 1-2); T₁-w GRE images (2.0 T) of phantom prior to (b), after (c) 47 and (d) 63 minutes of radiofrequency heating, inhomogeneous signal intensity in gel is due to air bubbles; (e) difference image after subtraction of (b) from (d). Note that the signal intensity from inserts 3 and 5 is almost unchanged after heating as the temperature never exceeded T_c.



FIG.4




FIG.5

5/13




FIG.6






FIG.7



7/13




FIG. 10




FIG. 11

8/13



FIG.12




FIG.13




Figure 14





Figure 15



**Figure 17**

12/13




Figure 18




Figure 19




Figure 20

Attorney Docket No.:

DECLARATION FOR PATENT APPLICATION AND APPOINTMENT OF ATTORNEY

As a below named inventor, I hereby declare that my residence, post office address and citizenship are as stated below next to my name: I believe that I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

USE OF PARTICULATE CONTRAST AGENTS IN DIAGNOSTIC IMAGING FOR STUDYING PHYSIOLOGICAL PARAMETERS

the specification of which (check one):

() is attached hereto, or () was filed on:

No.: and (if applicable) was amended on:

as U.S. Application Serial No. or PCT International Application

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment(s) referred to above. I acknowledge the duty to disclose information which is material to patentability as defined in *Title 37, Code of Federal Regulations, §1.56*. I hereby claim foreign priority benefits under *Title 35, United States Code, §119* of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed.

PRIOR FOREIGN APPLICATION(S)			PRIORITY CLAIMED	
Number	Country	Day/Month/Year Filed	Yes	No
9807840.5	GB	9 April 1998	YES	
9828874.9	GB	31 December 1998	YES	

I hereby claim the benefit under *Title 35, United States Code, §119(e)* of any U.S. provisional applications listed below:

Application Number	Day/Month/Year filed
60/119808	12 February 1999

I hereby claim the benefit under *Title 35, United States Code, §120* of any United States application(s) or PCT international application(s) designating The United States of America listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in that/those prior application(s) in the manner provided by the first paragraph of *Title 35, United States Code, §172*, I acknowledge the duty to disclose information which is material to patentability as defined in *Title 37, Code of Federal Regulations, §1.56* which became available between the filing date of the prior application(s) and the national or PCT international filing date of this application:

Application Number	Filing date	Status - Patented, Pending or Abandoned
PCT/G899/01100	9 April 1999	pending

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like are made are punishable by fine or imprisonment, or both, under *§1001 of the United States Code* and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

POWER OF ATTORNEY: I (We) hereby appoint as my(our) attorney, with full powers of substitution and revocation, to prosecute this application and transact all business in the Patent and Trademark Office in connection therewith: J. Ernest Kennay, Reg. No. 19,179; Eugene Mar, Reg. No. 25,893; Richard E. Fichter, Reg. No. 26,382; Charles R. Wolfe, Jr., Reg. No. 28,680; Thomas J. Moore, Reg. No. 28,974; Bruce H. Troxell, Reg. No. 28,592; Joseph DeBenedictis, Reg. No. 28,502; and

I (we) authorize my(our) attorneys to accept and follow instructions from Frank B. Dehn & Co. regarding any matter related to the preparation, examination, grant and maintenance of this application, any continuation, continuation-in-part or divisional based thereon, and any patent resulting therefrom, until I (we) or my(our) assigns withdraw this authorization in writing.

Send correspondence to: **BACON & THOMAS**
625 Sisters Lane - 4th Floor
Alexandria, VA 22314

Telephone calls to: **Richard E. Fichter**
(703) 683-0500

Full Name of First or Sole Inventor FOSSHEIM, Sigrid, Lise	Citizenship Norwegian
Residence Address Nycomed Imaging AS, Nycoveien 1-2, N-0401 Oslo, Norway	Post Office Address () Same as Residence
DATE 12/19/00	SIGNATURE S. Foss2

(x) See following page(s) for additional joint inventors.

Attorney Docket No. _____

DECLARATION FOR PATENT APPLICATION AND APPOINTMENT OF ATTORNEY

Page 2

Full Name of Joint Inventor KLAVENESS, Jo	Citizenship Norwegian
Residence Address Nycomed Imaging AS, Nycoveien 1-2, N-0401 Oslo, Norway	Post Office Address () Same as Residence
DATE 29/9/00	SIGNATURE <i>Jo Klavness</i>
Full Name of Joint Inventor BJORNERUD, Atle	Citizenship Norwegian
Residence Address Nycomed Imaging AS, Nycoveien 1-2, N-0401 Oslo, Norway	Post Office Address () Same as Residence
DATE	SIGNATURE
Full Name of Joint Inventor RONGVED, Pal	Citizenship Norwegian
Residence Address Nycomed Imaging AS, Nycoveien 1-2, N-0401 Oslo, Norway	Post Office Address () Same as Residence
DATE 29/9-00	SIGNATURE <i>Pal Rongved</i>
Full Name of Joint Inventor GOLMAN, Klaes	Citizenship Danish
Residence Address Nycomed Innovation AB, Ideon, S-205 12 Malmo, Sweden	Post Office Address () Same as Residence
DATE	SIGNATURE
Full Name of Joint Inventor SKURTVEIT, Roald	Citizenship Norwegian
Residence Address Nycomed Imaging AS, Nycoveien 1-2, N-0401 Oslo, Norway	Post Office Address () Same as Residence
DATE 2/10-00	SIGNATURE <i>Roald Skurtveit</i>

Attorney Docket No. _____

DECLARATION FOR PATENT APPLICATION AND APPOINTMENT OF ATTORNEY


Page 2

Full Name of Joint Inventor KLAIVENESS, Jo	Citizenship Norwegian
Residence Address Nycomed Imaging AS, Nycoveien 1-2, N-0401 Oslo, Norway	Post Office Address <i>(Same as Residence)</i>
DATE	SIGNATURE
Full Name of Joint Inventor BJORNERUD, Atle	Citizenship Norwegian
Residence Address Nycomed Imaging AS, Nycoveien 1-2, N-0401 Oslo, Norway	Post Office Address <i>(Same as Residence)</i>
DATE <i>Atle Bjørnerud 29/9/00</i>	SIGNATURE <i>Atle Bjørnerud</i>
Full Name of Joint Inventor RONGVED, Pal	Citizenship Norwegian
Residence Address Nycomed Imaging AS, Nycoveien 1-2, N-0401 Oslo, Norway	Post Office Address <i>(Same as Residence)</i>
DATE	SIGNATURE
Full Name of Joint Inventor GOLMAN, Klaes	Citizenship Danish
Residence Address Nycomed Innovation AB, Ideon, S-205 12 Malmö, Sweden	Post Office Address <i>(Same as Residence)</i>
DATE	SIGNATURE
Full Name of Joint Inventor SKURTVÆIT, Róald	Citizenship Norwegian
Residence Address Nycomed Imaging AS, Nycoveien 1-2, N-0401 Oslo, Norway	Post Office Address <i>(Same as Residence)</i>
DATE	SIGNATURE

Attorney Docket No: _____

DECLARATION FOR PATENT APPLICATION AND APPOINTMENT OF ATTORNEY

Page 2

Full Name of Joint Inventor KLAVENESS, Jo	Citizenship Norwegian
Residence Address Nycomed Imaging AS, Nycoveien 1-2, N-0401 Oslo, Norway	Post Office Address (/ Same as Residence)
DATE	SIGNATURE
Full Name of Joint Inventor BJORNERUD, Atle	Citizenship Norwegian
Residence Address Nycomed Imaging AS, Nycoveien 1-2, N-0401 Oslo, Norway	Post Office Address (/ Same as Residence)
DATE	SIGNATURE
Full Name of Joint Inventor RONGVED, Pal	Citizenship Norwegian
Residence Address Nycomed Imaging AS, Nycoveien 1-2, N-0401 Oslo, Norway	Post Office Address (/ Same as Residence)
DATE	SIGNATURE
Full Name of Joint Inventor GOLMAN, Klaes	Citizenship Danish
Residence Address Nycomed Innovation AB, Ideon, S-205 12 Malmo, Sweden	Post Office Address (/ Same as Residence)
DATE 21/9-02	SIGNATURE 
Full Name of Joint Inventor SKURTVEIT, Roald	Citizenship Norwegian
Residence Address Nycomed Imaging AS, Nycoveien 1-2, N-0401 Oslo, Norway	Post Office Address (/ Same as Residence)
DATE	SIGNATURE